



Ethnopharmacological values of Sacred Trees of Big Temples in Cuddalore district, Tamil Nadu, India

D. Sivalingam*, R. Rajendran and K. Anbarasan.

Department of Botany, Annamalai University, Annamalai Nagar-608 002.

Correspondence author: ssivlingam51@gmail.com

ABSTRACT

This study provides a significant ethnopharmacological informations, about the sacred plants of big temples in cuddalore district, tamilnadu state qualitatively and quantitatively. Ethanobotanical studies are very important to reveal the multi sort values of plants in the world in past and present. In this way since this is the first ethanobotanical study which records the traditional important medicinal values of sacred plants of big temples in cuddalore district. The information was obtained through open and semi-structured interviews with 35 (22 males, 13 females) knowledgeable local people and traditional healers. The collected data were analyzed qualitatively and quantitatively. In addition, conservation status of the plants, usage value and relative importance were determined.

*A total of six species of sacred plants, mostly trees, belonging to six families were studied in this investigation. They were used to treat 30 diseases and 12 major ailment categories. Leaves were the most frequently used plant part. Based on IUCN red data, the identified sacred plants include one least concerned species, one vulnerable species and two endangered and threatened species. The ailment categories of Gastro-intestinal problems had the highest percentage (83.3) of use plant species (Table-5). The most important species according to their high use value (UV) were *Aegle marmelos* (0.97) and *Prosopis cineraria* (0.94) which are the most frequently and popularly used plants. The species which had the highest Relative Importance (RI) values are *Aegle marmelos* (1.80) and *Ficus religiosa* (1.71). As a result of the present study, it can be concluded the sacred plants play an important role in the health care of human being and people rely on medicinal plants to treat various ailments and diseases. In addition, the ethnomedicinal plants with high UV and RI values might give some useful leads for further pharmacological investigation.*

Key words: Sacred trees, Conservation, Ethnopharmacology, Ethano-medicine, Sthalavriksha, Big Temple Cuddalore district.

Received 10.12.2015

Revised 25.12.2015

Accepted 10.01.2016

INTRODUCTION

Plants are the important source of therapeutic drugs and play a significant role in the survival of many tribal and ethnic communities [1]. According to World Health Organization (WHO), 80% of the developing countries rely on the traditional medicines and in which 85% of use plants or their extracts are used as plant drugs for their healthcare need [2,3]. India is one of the treasure houses of medicinal plants in the world. The survey of medicinal plants in a particular area is important to conserve the traditionally important plants of that particular landscape [4].

Sthalavriksha is referred to as a plant which is venerated from time immemorial by devotees as being as holy as the presiding deity of a temple [5]. The sanskrit term sthalavriksha means tree of the locality (sthal-place; vriksha-tree). Sthalavriksha is a natural tree found in the temple site before construction of the temple and most temple myths (Sthalapuranas) and temple histories (Sthalavaralaru) refer to a prime diety that was first unearthed or found under the tree [6]. Due to traditional beliefs, both the devotees and temple authorities serve as protectors of the sthalavriksha in temples and if a tree (sthalavriksha) dies because of old age, it is usually replaced by a sapling from the same species. Hence, the tree occurs constantly in a temple for several centuries. However, in most cases the original sthalavriksha is still living within the temple grounds. In Hindusim, especially in Shaivism, there are three important aspects of the temple grounds, Moorthy (a Deity), Sthalam, (a Shrine and Sthalavriksha) and a Theertham, (Sacred tank or water body). These are the three prime elements to learn about the antiquity of a temple. The worship of these three elements will yield wisdom even without a guru or teacher [7]. Sthalavriksha worship is mostly associated with Shaivism (worship of Lord Siva), Vaishnavam (worship of Lord Vishnu)

and Sthalavriksha gets divine power from these deities, which are treated as equal to the prime deity of the temple [9]. Many medieval Tamil sacred hymns, e.g., Devaram [8] and Thiviyaprabantham [10] refer to Sthalavrikshas and their associated deities. Even though Sthalavriksha worship is an ancient practice in Tamil Nadu a very few studies have been conducted on its ethno importance [11, 12]. In particular, ethnomedicinal uses of Sthalavriksha were referred to, based on secondary sources only. Hence, the present study was designed to gather data using a field survey with special reference to: 1. To catalogue all Sthalavriksha species and their associated deities in the temples of cuddalore district, Tamil Nadu. 2. To document the ethnomedicinal utilization of user groups including devotees, priests, Nattuvaidiyas on Sthalavriksha and Executive officers of the temples studied.

MATERIALS AND METHODS

Study area

Cuddalore districts is located (11°43' N and 79°49' E) in the East Coast about 23 kms south of Puducherry region. Various types of soil are found in the Cuddalore region which include red loamy, coastal alluvium, delta alluvium, red laterite, deep black and red sandy. The mean annual rain fall is 1,079mm and the dry season lasts for six months (January to June), and receives less than 60mm rainfall on monthly average. The mean annual minimum and maximum temperature are 22.75°C and 33.64°C, respectively.

Field trips were made to collect information's about six sacred trees of big temples in cuddalore district, namely i) *Ficus religiosa* (L.) (Moraceae), ii) *Stereospermum suaveolens* D.C. (Bignoniaceae), iii) *Aegle marmelos* (L.) (Rutaceae), iv) *Prosopis cineraria* (L.) (Mimosaceae), v) *Exoecaria agallocha* (L.) (Euphorbiaceae) and vi) *Pterocarpus marsupium* (L.) (Fabaceae), informations obtained from Bhuvarahaswami Thirukovil, Srimushnam; Padaleshwarar Temple, Cuddalore; Veeranateswarar Temple, Panruti; Vridhagireeswarar Temple, Vriddhachalam; Natarajar Temple, Chidambaram and Vaidyanathasami Temple of Tittagudi respectively.

The local people including village heads, traditional healers and traders of cuddalore district who have indigenous knowledge about sacred plants of their locality and informations were collected by group discussions and interviews with them in their local language (Tamil). Binomials of the plants with family, their local name(s), part(s) used and therapeutic uses were recorded. Voucher specimens were collected and identified with the standard floras [29, 30]. All the voucher specimens were maintained in the herbarium centre of Department of Botany, Annamalai University, Chidambaram (India).

Interview and ethnomedicinal data collection

The ethnomedicinal information was collected using questionnaires by semi-structured interviews. A total of 35 informants (22 males and 13 females) include various strata of participants (local users, knowledgeable persons, village heads and traditional vaidyas/healers with the age ranged between 40-75 years) were selected for interview and get the data.

Collection of plants

Generally, the plant specimens were collected with flower and fruit conditions. In case, if there is no flower and fruit conditions, the plant twig with few leaves were collected for proper identification and herbarium preparation. For herbarium preparation, standard procedures given by [18].

Taxonomic identification

The sacred plants were identified and their scientific names were known with the help of professional/subject experts and the reputed flora and books [13-17, 19].

Conservation status of the present sacred plants

In this study, collected sacred species were compared with IUCN Red List in order to identify their living status [20].

Data analysis

The majority of the data collected in this study were descriptive in nature. Interview data were coded and sorted into themes. Inconsistencies and unique statements were noted and given particular attention. Recurrent themes were uncovered in this report by a process of systematic content analysis. In its broadest sense, different researchers have emphasized various aspects of content analysis, from its capacity to generate quantitative descriptions by analyzing work counts [21]. Ethnopharmacological data were analyzed and summarized by using Microsoft excel.

Use value (UV)

The use value (UV) [21], a quantitative method that demonstrates the relative importance of species known locally was also calculated using the formula $UV = (\sum U/n)$.

Where UV is the use value of species, 'U' is the total number of use reports per species and 'n' represents the total number of informants interviewed for a given plant. Values will be high (near 1) if there are

many use reports for a plant, implying that the plant is important, and near 0 if there are few reports related to its use.

Relative importance (RI)

Relative importance value (RI) was calculated according to Bennett and Prance [22]. $RI = PP + AC$, where PP is obtained by dividing the number of pharmacological properties (reported by dividing the number of pharmacological properties (reported specific ailments) attributed to a species divided by the maximum number of properties attributed to the most resourceful species (species with the highest number of properties). AC is the number of ailment categories treated by a given species divided by the maximum number of ailment categories treated by the most resourceful species. The highest possible value of RI is 2.0, which indicates the highest diversity of medicinal uses of a plant [23].

RESULTS AND DISCUSSION

Conservation status of the plants

According to IUCN Red List Categories and Criteria (Version 3.1), four types of species are found, Vulnerable (Vu), Endangered (En), Threatened (T) and Least concerned (LC) at local level. For the present species it is given in table 3.

Use value (UV)

Use value representing the relative importance of plants, were high for *Aegle marmelos* (0.97) and *Prosopis cineraria* (0.94). The lowest use value was calculated at 0.80 (Table- 4). High use-value plants were the most frequently used sacred plant species.

Relative importance

The plant with more number of pharmacological properties (PP) were *Aegle marmelos*, *Prosopis cineraria* (7 PP); so they have a normalized PP value of 1.00 (7/7). *Ficus religiosa* was employed to treat five ailments categories and had a normalized AC value of 1.00 (5/5). *Aegle marmelos* had the highest RI of 1.80 and it was followed by *Ficus religiosa* (1.71) (Table-6).

Aegle marmelos L.

Family – Rutaceae, Local Name - Bel; Sanskrit-Sriphla, Bilva; English- the Bengal Quince or the Bel tree.

Sacred value: Plant is generally grown in temples and it is sacrilege to cut it down. The tree is sacred to the Trimurti (a representative of Shiva) and Parvati (wife of Lord Shiva). The leaves are used in enchantments and twigs are used for sacred fire during Samidhas. Leaves and fruits are used in the worship of Lord Shiva.

Medicinal value: Fruit's fleshy part is dried, powdered and given to children as anastigmatic for diarrhoea, while fruits and pulp is used for stomach ache. Juice is prepared from ripped fruit, used as mild laxative. Leaves chewed and swallowed at every morning help in healing stomach ulcer and also to reduce sugar in diabetes patient.

Ficus religiosa L.

Family- Moraceae, Local name - Peepal; Sanskrit-Aswaththama; English- Bo-tree, Peepal

Sacred values: The tree is sacred for Hindus, viewed as the female to the Banyan tree. He further says that it is the transformation of the Gods Guru- Ashwath. The tree is worshipped on every month on which a new moon falls. Among the hindus five sacred trees (i.e. Peepal, Gular, Bargad, Pakar, and Mango) Peepal ranks first. A good hindu (Sanskari Hindu) while starting his journey if he sees a peepal tree, then he repeats the verse which may be translated - the roots Brahma, the bark Vishnu, and the branches Mahadeos. The Peepal is believed to be inhabited by the sacred triad – Brahma, Vishnu and Mahesh (Shiva) of its wood the spoons are made with which to pour ghee on the sacred fire (Home).

Medicinal value: Bark paste applied to throat of children (3-4 times a day) to get relief from apathies. Leaf bud is used in curing snake bite. Aqueous extract of bark shows anti-bacterial activity.

Exoecaria agallocha (Euphorbiaceae) is known as 'blind-your-eye' is a typical mangrove associated species that occurs along the coastal areas of Tamil Nadu State, India. Its common name refers to the damaging effect of its white milky sap which causes blindness in human when have on human eyes.

Sacred values: This tree has religious significance for the local community and people who visit the Lord Natarajar Temple in Chidambaram town revere this sacred plant.

Medicinal value: Various parts of this plant have been used in the traditional medicine for the treatment of ulcers, leprosy and also as an aphrodisiac for decades [16]. In Sri Lanka, the smoke of the burning wood has been used in the treatment of leprosy, while the root pounded with ginger has been used to reduce swellings of hands and feet [24]. A novel Phorbol ester, an anti-HIV principle has also been isolated from the leaves and stem of this unique plant [15]. Fatty acids are widely occurring in natural fats and dietary oils and they play an important role as nutritious substances and metabolites in living organisms [25]. Many fatty acids are known to have antibacterial and antifungal properties [26]. However, little is known on the antibacterial and antifungal properties of *Exoecaria agallocha*.

Pterocarpus marsupium (Fabaceae); Common name: Indian Kino, Gummy Kino, Marsupium.

Sacred values: Leaves and Flowers are offered to God for pooja.

Chemical constituents and components: Main chemical components are liquiritigenin, isoliquiritigenin, pterosupin, epicatechin, pterostilbene, kinotannic acid, beta-eudesmol, marsupol, carpusin, marsupinol, kinoin and kinored.

Actions- Pterostilbene: 1. It lowers the lipid and glucose level in body. 2. It shows significant reduction of glycosylated haemoglobin and an increase in total haemoglobin level. 3. It inhibits cell proliferation and induces apoptotic effect in tumor cell lines. 4. It shows moderate inhibition of cyclo oxygenase (cox)-1 and weakly active against cox-2, thus shows potent antioxidant activity. 5. It inhibits electrolyte leakage caused by herbicide-induced oxidative damage.

Alkaloids: It significantly lowers the high blood sugar level in body.

Tannins: 1. It shows scavenging activity against mitochondrial lipid peroxidation. 2. It causes significant decrease in cholesterol level. 3. It shows antimicrobial activity against bacteria and virus.

Curing Diseases: 1. It is useful in diabetes and heart problems. 2. It is helpful in controlling skin diseases. 3. It is also beneficial in the treatment of fractures, bruises, leprosy, leucoderma, constipation, depurative, rectalgia, ophthalmopathy, hemorrhages and rheumatoid arthritis [28].

Stereospermum suavelons (Bignoniaceae) is a medicinal tree species native to India [31]. The Bignoniaceae having about 100 genera with 800 species.

Sacred value: Sthalavriksham of Padaleshwarar Temple, to fulfil the vow of getting a child, yellow cloth cradles are tied in the branches of this tree by childless women (Fig.1).

Medicinal value: It is known for their antimicrobial, antiprotozoal, and anti-inflammatory properties [32, 33]. Both the timber [34] and the root heartwood [35] of *S. suaveolens* were found to contain lapachol, elicitor of contact dermatitis [36]. Moreover, barks, flowers, roots and leaves of *S. suaveolens* are used by traditional healers, rural communities and pharmaceutical companies for remedies of diseases like heating, vomiting, eructation, piles, acidity, diarrhoea, gonorrhoea, loss of taste, malaria and other fevers [31].

Prosopis cineraria (Mimosaceae)

The tree is found in extremely arid conditions, with rainfall as low as 150 mm annually; but is indicative of the presence of a deep water table. It has high tolerance to high alkaline and saline environments [37].

Social value: The wood of *P. cineraria* is a good fuel source, and provides excellent charcoal. The leaves and pods are consumed by livestock and are beneficial forage. In Rajasthan, India, *P. cineraria* is grown in an agro forestry setting in conjunction with millet. The tree is well suited for an agro forestry setting, because it has a single-layered canopy, it is a nitrogen fixer (thus enriching the soil), and its deep roots avoid competition for water with crops [38].

Medicinal value: Gives relief from muscular and joint pain and snake poison and also acts as anti-inflammatory and flatulence. (Tables 1 & 2).

Table 1: Particulars about Sacred Trees (Sthalavrikshas) studied.

S.No.	Name of the Sacred Tree (Sthalavriksha)	Family	Vernacular Name	Name of the Temple and place	Deity of the temple
1.	<i>Aegle marmelos</i> L.	Rutaceae	Vilvam	Veeranateswarar Temple, Tiruvatikai, Panruti	Shivan
2.	<i>Exoecaria agallocha</i> L.	Euphorbiaceae	Thillai	Natarajar Temple, Chidambaram	Shivan
3.	<i>Ficus religiosa</i> L.	Moraceae	Arasu	Bhuvanahaswami Temple, Srimushnam	Shivan
4.	<i>Prosopis cineraria</i> L.	Mimosaceae	Vanni	Vridhagireeswarar Temple, Vriddhachalam	Shivan
5.	<i>Pterocarpus marsupium</i> Roxb.	Fabaceae	Vengai	Vaidyanathaswami Temple, Tittagudi	Shivan
6.	<i>Stereospermum suavelons</i> DC.	Bignoniaceae	Paathiri	Padaleshwarar Temple, Cuddalore	Shivan

Table 2: Sacred trees and their religious and medicinal uses.

S.No	Botanical name	Religious uses and belives	Medicinal uses
1.	<i>Aegle marmelos</i> L. Correa	i) Sthalavriksham of Veeranateswarar Temple ii) Leaves and flowers are offered to God for pooja.	i) 25g of root powder is boiled in water and filtered decoction is taken thrice a day for 7 days to cure fever. ii) Half of ripe fruit is taken twice a day for 3-4 days to cure constipation and dyspepsia.
2.	<i>Exoecaria agallocha</i> L.	i) Sthalavriksham of Natarajar temple.	i) The roots are used to treat toothache and swellings ii) A novel phorbol ester, an anti-HIV principle has also been isolated from the leaves and stem of this unique plant.
3.	<i>Ficus religiosa</i> L.	i) Sthalavriksham of Bhuvarama Swami Temple. ii) Male offspring is entreated, if pious woman move around its trunk 108 times.	i) Milky latex is applied externally to cure foot cracks, and healing of blood clotted site. ii) 20-50g of root bark is made in to a powder and one tea spoonful of powder is mixed with little amount of coconut oil and is applied externally on the blood clotted site to cure blood clotting.
4.	<i>Prosopis cineraria</i> L.	i) Sthalavriksham of Vridhagireeswarar Temple ii) To fulfil the vow of getting a child, yellow cloth cradles are tied in the branches of this tree by childless women	i) Gives relief from muscular and joint pain and snake poison and also acts as anti-inflammatory and flatulence.
5.	<i>Pterocarpus marsupium</i> Roxb.	i) Sthalavriksham of Vaidyanatha Swami Temple	i) It is useful in diabetes and heart problems. ii) It is helpful in controlling skin diseases. iii) It causes significant decrease in cholestral level. iv) It shows antimicrobial activity against bacteria and virus.
6.	<i>Stereospermum suavelons</i> DC	Sthalavriksham of Padaleswarar Temple	Flowers pounded with honey are taken for highcough.

Table 3: IUCN Red listed Sacred plants recorded in study area.

S.No	Botanical Names	Family	IUCN status
1.	<i>Aegle marmelos</i> L.	Rutaceae	T
2.	<i>Exoecaria agallocha</i> L.	Euphorbiaceae	En
3.	<i>Ficus religiosa</i> L.	Moraceae	LC
4.	<i>Prosopis cineraria</i> L.	Mimosaceae	T
5.	<i>Pterocarpus marsupium</i> Roxb.	Fabaceae	Vu
6.	<i>Stereospermum suavelons</i> DC.	Bignoniaceae	En

LC: Least concerned, T: Threatened, En: Endangered, Vu: Vulnerable

Table 4: Sacred plants – Ethanomedicinal value and use value (UV).

S.No	Botanical Names	Parts used	Disease Treated	UV value
1.	<i>Aegle marmelos</i> L. (Rutaceae)	Leaf paste Leaves Fruit pulp Fruit pulp with milk Leaves (sweet leaves) Leaves & Fruit Leaf juice Leaf juice & Fruit paste	Knee and joint pain, infertility Blood sugar reduction Skin boils Diarrhea Diabetes, skin disease Cold and cough Menstrual disorder in women Cold, cough, chest diseases	0.97
2.	<i>Prosopis cineraria</i> L. Druce (Mimosaceae)	Bark Leaves Flowers Gum	Tooth ache Cold, cough, fever Prevent miscarriage Dysentery	0.94

3.	<i>Excoecaria agallocha</i> L. (Euphorbiaceae)	Flowers Root Oil	Eye diseases Tooth ache, intestinal worms Joint pain, leprosy	0.8
4.	<i>Pterocarpus marsupium</i> Roxb. (Fabaceae)	Bark Flowers Gum	Tooth ache Fever Tooth ache	0.85
5.	<i>Ficus religiosa</i> L. (Moraceae)	Bark Fruit Latex Leaves Seeds	Ulcer Laxative Piles, diarrhea Cut wounds Refrigerant	0.88
6.	<i>Stereospermum suavelons</i> DC. (Bignoniaceae)	Flowers	Diabetic boils	0.91

Table 5: Major categories of ailments

Ailment categories	Biomedical terms	No. of species used	Percentage
Gastro-intestinal problems	Intestinal problems, Stomachic, anthelmintic, diarrhea, dysentery, cholera, carminative, indigestion, dyspepsia, vermifuge, hiccup, flatulence, laxative, purgative, antispasmodic, appetizer, deobstruent, cathartic, colicpain, anorexia, nausea, antibilious	5	83.3
Respiratory problems	Cough, bronchitis, asthma, hemoptysis, expectorant, tuberculosis, scrofula	3	50.0
Urinary and rectal problems	Hematuria, piles, dysuria, kidney, urinary, urethrorhea, nephritis, crystalluria, lithontriptic, strangury, calculi, constipation	2	33.3
Circulatory diseases	Hypertension, anemia, styptic, astringent, blood purifier, hemorrhage, dropsy, depurative	1	16.6
Infections and parasitic diseases	Elephantiasis, antiseptic, ascariasis, chicken pox, head lice, ringworm, scabies, amebiasis, antiprotozoal	2	33.3
Inflammations and pains	Abdominal pain, rheumatism, narcotic, anodyne, abdominal pain, sedative, pectoral pain, cephalalgia, fever, analgesic, antipyretic, abdominal pain, febrifuge	4	66.6
Dermatological problems	Leprosy, rubefacient, antiscorbutic, demulcent, cooling, eczema, leukodermatic, emollient, diaphoretic, eruptions, psoriasis, erysipelas, dermatitis, boils, skin diseases, suppurative	2	33.3
Female problems	Emmenagogue, galactagogue, menorrhagia, leucorrhoea, dysmenorrhoea, hemorrhage, vaginal laxity, vaginal disinfectant, abortifacient	2	33.3
Male fertility problems	Seminal weakness, sperm coagulant, spermatorrhoea, Aphrodisiac	1	16.6
Eye, ear, nose and throat problems	Gout, otalgia, ophthalmia, pharyngitis, rhinitis, cold, nyctalopia	3	50
Endocrine, nutritional and metabolic disorders	Stimulant, alterative, beriberi, diabetes, scurvy, splenomegaly, diuretic	2	33.3
Injury and poisons of external causes	Snake bite, allergy, burns, cuts, wounds, fracture	1	16.6

Table 6: Relative Importance (RI) values of medicinal plants used against five specific use categories and seven ailments categories treated.

Plant species	PP ^a	AC ^b	RI ^c
<i>Aegle marmelos</i> L.	1.00	0.80	1.80
<i>Excoecaria agallocha</i> L.	0.28	0.20	0.48
<i>Ficus religiosa</i> L.	0.71	1.00	1.71
<i>Prosopis cineraria</i> L.	1.00	0.60	1.60
<i>Pterocarpus marsupium</i> Roxb.	0.85	0.80	1.65
<i>Stereospermum suavelons</i> DC.	0.42	0.40	0.82

^aPP : Pharmacological properties
^bAC: Ailment categories
^cRI : Relative importance (PP+AC)

Fig. 1 - Padaleshwarar Temple (Cuddalore) Sthalavriksham - *Stereospermum suavelons* DC



CONCLUSION

Sthalavrikshas are valued for their botanical, medicinal, environmental, religious and mythical importance. The sthalavrikshas of Tamilnadu constitute a part of genetic resources for the conservation of species diversity. Propagation of sthalavrikshas in temples contributes to the conservation of our floral diversity. Some trees are important for their economic role in ship building or in the timber industry, some for providing homes for various animals, birds and others for their medicinal and air purifying qualities. In the present study, it is concluded that the religious activities are having close relationship with plants boost up the mental health of local people of Cuddalore district and many of the sacred plants found in the household and temples were used for various religious cultural activities as well as for health care. These sacred plants are worshiped by the local people for getting the blessing of health and wealth by positive powers of nature. Hence the religious ceremonies, rites act as a protective factor or device for the conservation of sacred plants. So, it is the duty of present generation to preserve and promote these aesthetic treasures to conserve biodiversity and nature, which will surely play a part in progeression of human beings. These sacred trees preserved through millennia by our ancestors as potential bio resources should be respected and conserved for the future generation.

ACKNOWLEDGMENTS

The authors are cordially grateful to the people inhabiting the study area for sharing their traditional knowledge and also thankful to the Department of Botany, Annamalai University. The authors gratefully acknowledge the University Grants Commission (UGC), New Delhi, for the financial support in the form of BSR meritorious Fellowship. We specially express our sincere thaks to Mr. R. Rajendran, for providing ethanomedical information about the study plants.

REFERENCES

1. Francis Xavier, T., Kannan, M., Auxilia, A., (2015). Observation on the traditional phytotherapy among the Malayali tribes in Eastern Ghats of Tamilnadu, South India. *Journal of Ethnopharmacology* 165 (2015) 198 – 214.
2. Senthilkumar, K., Aravindhan, V., Rajendran, A., (2013). Ethnobotanical survey of medicinal plants used by Malayali tribes in Yercaud Hills of Eastern Ghats, India. *Journal of Natural Remedies* 13 (2), 118–132.
3. Sheldon, L.M., Balick, M., Laird, S.A., (1998). Is using medicinal plants compatible with conservation? *Plant Talk* 98, 29 – 31.
4. Sivasankari, B., Anandharaj, M., Gunasekaran, P., (2014). An ethnobotanical study of indigenous knowledge on medicinal plants used by the village peoples of thoppampatti, Dindigul district, Tamilnadu, India. *Jornal of Ethnopharmacology* 153 ,408 – 423.
5. Gunasekaran, M., and Balasubramanian, P., (2005). Sthalavriksha worship; A tool in plant conservation in Tamil Nadu. Pp. 163-166 in *Proceedings of National Strategy for Conservation of Sacred Groves*. Edited by C. Kunhikannan & G. Singh. Institute of Forest Genetics and Tree Breeding, Coimbatore, India.

6. Gunasekaran, M., and Balasubramanian, P., (2012). Sthalavriksha worship; A tool in plant conservation in Tamil Nadu. Pp. 163-166 in Proceedings of National Strategy for Conservation of Sacred Groves. Edited by C. Kunhikannan & G. Singh. Institute of Forest Genetics and Tree Breeding, Coimbatore, India.
7. Thambiran, M., (1963). Thayumanavar's Paraparkkani. Kasimadam, Thirupananthal, India.
8. Thambiran, M., (1997). Thirugnanasambathar Devaram, Volume1-6. Dharumapuram Adeenam, Myladuthurai, India.
9. Nedunchezhiyan, (2005). Thamizhar Kanda Thavaraviyal. International Institute of Tamil Studies, Chennai, India.
10. Anonymous, (1962). Nalayirathiviyaprapantham. Madavadasan, Ramakrishana Mudaliyar, Madras, India.
11. Amirthalingam, M., (1998). Sacred Trees of Tamil Nadu; A survey. C.P.R. Environmental Education Centre, Chennai, India.
12. Thirugnanam, S., (1995). Thirukoil Marangalin Maruthuvapayangal. Selvi Pathipagam, Tiruchi, India.
13. Hooker, J.D., 1884. The Flora of British India. L. Reeve and Co. Kent.
14. Hooker, J.D., 1973. The Flora of British India, vol. I-VII, pp. 1872-1897 (Reprinted by Bishen Singh Mahendra Pal Singh, Dehradun and Periodica IExperts, Delhi).
15. Karalai, C., Wiriyachitra, P., Opferkuch, H. J., Hecker, E. 1994. Cryptic and free skin irritants of the daphnane and tiglane types in latex of *Excoecaria agallocha*. *Planta Medica*, 60, 351-355.
16. Kirtikar, K. R., Basu, B.D., (1999). Indian Medicinal Plants. Vol. IIV, Lalit Mohan Basu Publishers, Allahabad, India.
17. Kritkar, K.R., Basu, B.D., (1975). Indian Medicinal Plants, vol. I-IV. Periodical Experts, Delhi, India.
18. Jain, S.K., Rao, R.R., 1977. A Handbook of Field and Herbarium Methods. Today and Tomorrow's Tomorrow's Printers and Publishers, New Delhi
19. Jain, S.K., De Filippis, R.A., 1991. *Medicina Plant of India*, vol. I-II. Reference Publications Inc., Algonac, MI.
20. IUCN, 2013. IUCN Red list of Threatened species, Version 2013.1. (www.iucnredlist.org).
21. Silverman, D., 1993. *Interpreting Qualitative Data: Methods for Analyzing Talk, Text and Interaction*. Sage Publications, Thousand Oaks, California
22. Trotter, R.T., Logan, M.H., 1986. Informant consensus: a new approach for identifying potentially effective medicinal plants. In: Etkin, N.L. (Ed.), *Plants in Indigenous Medicine and Diet, Behavioural Approaches*. Redgrave Publishing Company, Bredford Hills, New York, pp. 91-112.
23. Sivasankari, B., Pitchaimani, S., Anandharaj, M., (2013). A study on traditional medicinal plants of Uthapuram, Madurai District, Tamilnadu, SouthIndia. *Asian Pac. J. Trop. Biomed.* 3 (12), 975-979.
24. Bennett, B.C., Prance, G.T., (2000). Introduced plants in the indigenous pharmacopoeia of northern South America. *Econ. Bot.* 54, 90-102.
25. Jayaweera, D.M.A., (1980). Medicinal plants used in Ceylon. *J. Natl. Sci. Coun. Sri Lanka*, 2, 214-215.
26. Cakir, A., 2004. Essential oil and fatty acid composition of the fruits of *Hippophae rhamnoides* L. (Sea Buckthorn) and *Myrtus communis* L. from Turkey. *Biochem. System. Ecol.*, 32, 809-816.
27. Russel, A.D., (1991). Mechanisms of bacterial resistance to non antibiotics: food additives and food pharmaceutical preservatives. *J. Appl. Bacteriol.*, 71, 191-201.
28. Jahromi, M.A., and Ray, A.B., 1993. Antihyperlipidemic effect of flavonoids from *Pterocarpus marsupium*, *JNat Prod.* Jul; 56 (7): 989-994.
29. Gamble, J.S., 1936. *Flora of the Presidency of Madras. Vol I-III*. Allard & Co. London. (Reprinted - 1956) Botanical Survey of India. Calcutta.
30. Matthew, K.M., 1983. *The Flora of Tamilnadu Carnatic. The Rapinact Herbarium*, Tiruchirappalli, Tamilnadu.
31. Troup, R.S., 1986. *Silviculture of Indian Trees Volume 2: Leguminosae (Caesalpinieae) to Verbenaceae*. International Book Distributors, Dehradun, India.
32. Binutu, O.A., Adesogan, K.E., Okogun, J.I., 1996. Antibacterial and antifungal compounds from *Kigeliapinnata*. *Planta Med*, 62 (4): 352-353.
33. Onegi, B., Kraft, C., Köhler, I., Freund, M., Jenett-Siems, K., Siems, K., Beyer, G., Melzig, M.F., Bienzle, U., Eich, E., 2002. Antiplasmodial activity of naphthoquinones and one anthraquinone from *Stereospermum kunthianum*. *Phytochemistry*, 60 (1): 39-44.
34. Sandermann, W., Dietrichs, H.H., (1957). Untersuchungen über termitenresistente Holzarten. *Holz als Roh- und Werkstoff*, 15 (7): 281-297.
35. Joshi, K.C., Bansal, R.K., Patni, R., (1977). Chemical examination of the roots of *Stereospermum suaveolens* DC. *J Indian Chem Soc*, 54: 648-649.
36. Schulz, K.H., Garbe, I., Hausen, B.M., Simatupang, M.H., 1977. The sensitizing capacity of naturally occurring quinones. I. Naphtho quinones and related compounds. *Arch Dermatol Forsch*, 258: 41-52.
37. Tropical Forages, 2012. Centre for International Agriculture Research and Food and Agriculture Agency. 2005. Retrieved 2012-03-21.
38. World Agro forestry Centre. Retrieved, 2012-03-21.

CITATION OF THIS ARTICLE

D. Sivalingam, R. Rajendran and K. Anbarasan. Ethnopharmacological values of Sacred Trees of Big Temples in Cuddalore district, Tamil Nadu, India. *Bull. Env. Pharmacol. Life Sci.*, Vol 5 [3] February 2016: 39-46