



Relationship and Influence of Independent Variables On How To Do Knowledge Under Rainfed Condition-A Critical Analysis

B. Sudhakar* and K.A. Ponnusamy**

*Assistant Professor, Dept. of Agrl. Extension, Faculty of Agriculture, Annamalai University

**Retired Professor, Tamil Nadu Agriculture University (TNAU), Coimbatore.

ABSTRACT

The cotton is the major commercial crop in the world. The cotton production in India is was low as compared to many other countries. Under the impact of green revolution, due to monocropping, there have been many outbreaks of insect pests especially on rice and cotton. To meet these challenges, it is of utmost importance that in future the insect problems would have to be tackled through Integrated Pest Management (IPM). IPM has been defined by Pretty et al (1992) as the integrated use of some or all the pest control strategies in a way that not only reduce pest population to economically acceptable levels but it is sustainable and non-polluting. The IPM programme aims at educating the farmers and extension agencies through Farmers Field Schools (FFS). Coimbatore district stands first in total number of IPM-FFS training programmes conducted for cotton throughout the Tamil Nadu State over the years and hence, it was selected for the study. The highest area under cotton and maximum number of IPM -FFS training programmes conducted were considered as the criteria to select the Taluk representing rainfed condition. The same criteria were used for selection of Block where Avinashi block under rainfed condition were selected. In Avinashi block, four villages were selected. A sample of 100 farmers was selected for study. This part deals with the specific objective was to study the relationship and influence of profile characteristics with how to do knowledge of IPM oriented cotton growers under rainfed condition. When all the other variables are kept at constant level, a unit increase in educational status, mass media exposure and innovativeness increase in how to do knowledge by 1.3288, 0.331 and 3.0458 units respectively among IPM oriented cotton growers under rainfed condition.

Received 14.12.2018

Revised 19.01.2019

Accepted 29.01.2019

INTRODUCTION

The white gold occupied major parts of commercial crops. The area under cotton in India is the largest and constitutes nearly one-fourth of the World's cotton area. Under the impact of green revolution, due to monocropping, there have been many outbreaks of insect pests especially on rice and cotton. To meet these challenges, it is of utmost importance that in future the insect problems would have to be tackled through Integrated Pest Management (IPM). IPM has been defined by Pretty *et al* [4] as the integrated use of some or all the pest control strategies in a way that not only reduce pest population to economically acceptable levels but it is sustainable and non-polluting. The IPM programme aims at educating the farmers and extension agencies through Farmers Field Schools (FFS). Under FFS programme, farmers are made experts in identifying natural enemies of pests, monitoring regular pests and taking suitable management measures. In the year 1999-2000 under ICDP (Intensive Cotton Development Programme) totally 1500 FFS were organized and 45000 cotton growers were trained throughout India [1-4]. Coimbatore district stands first in total number of IPM-FFS training programmes conducted for cotton throughout the Tamil Nadu State over the years and hence, it was selected for the study.

SPECIFIC OBJECTIVE OF THE STUDY

The specific objective of this study was relationship and influence of how to do knowledge of Integrated Pest Management (IPM) adopting cotton growers with their profile characteristics under rainfed agro-ecosystem in Coimbatore district of Tamil Nadu, India.

MATERIAL AND METHODS

Coimbatore district stands first in total number of IPM-FFS training programmes conducted for cotton throughout the Tamil Nadu State over the years and hence, it was selected for the study. The highest area under cotton and maximum number of IPM -FFS training programmes conducted were considered as the criteria to select the Taluk representing rainfed condition. The same criteria were used for selection of Block where Avinashi block under rainfed condition were selected. In Avinashi block, four villages were selected. A sample of 100 farmers was selected for study. This part deals with the specific objective was to study the how to do knowledge of IPM oriented cotton growers under rainfed condition. How to do knowledge consists of information that is necessary to use an innovation properly. The adopter must understand what quantity of an innovation to secure, how to do use it correctly and so on [6]. The list of items that would help to measure the how to do knowledge on recommended IPM technologies were prepared in consultation with entomologists, extension scientists and by referring to the IPM-FFS guide.

RESULT AND DISCUSSION

1. Relationship of independent variables with how to do knowledge of IPM practices under rainfed condition

It may be seen from Table 1 that out of the sixteen variables studied under rainfed condition, the correlation coefficient value of nine variables viz., educational status, area under cotton cultivation, annual income, IPM equipment possession, extension agency contact, mass media exposure, innovativeness, risk orientation and economic motivation were found to have positive and significant relationship at 1 % level with how to do knowledge. The risk orientation had showed positive and significant relationship at 5 % level with awareness knowledge, whereas farm size showed negatively significant relationship at 1 % level with how to do knowledge.

The remaining characteristics viz., age, occupation, farming experience, experience in cotton cultivation, social participation and attributes of IPM were non-significantly correlated with how to do knowledge.

2. Influence of independent variables with how to do knowledge of IPM practices under rainfed condition

The results of multiple regression are given in Table 1, which indicated that out of the selected sixteen characteristics under rainfed condition, only three characters viz., educational status, mass media exposure and innovativeness had positive and significant influence towards the how to do knowledge among IPM oriented cotton growers under rainfed condition. The R² value indicated that, all the 16 characteristics taken together explained as much as 50.40 per cent of the variation in the how to do knowledge, *ceteris paribus*.

It may be inferred that when all the other variables are kept at constant level, a unit increase in educational status, mass media exposure and innovativeness increase in how to do knowledge by 1.076, 0.331 and 3.147 units respectively among IPM oriented cotton growers under rainfed condition.

Table1. Correlation and Multiple regression analysis of independent variables on how to do knowledge of IPM practices under rainfed condition

V.NO	Variables	'r' value	't' value	B value	Standard error
1	Age	-0.1086 ^{NS}	0.7932 ^{NS}	0.0981	0.1237
2	Educational status	0.5359**	3.1310**	1.3288	0.4244
3	Occupation	0.1232 ^{NS}	0.5188 ^{NS}	1.0996	2.1198
4	Farm size	-0.4080**	-0.2506 ^{NS}	-0.1677	0.6691
5	Area under cotton cultivation	0.3143**	-1.1082 ^{NS}	-0.8611	0.7770
6	Farming experience	-0.1836 ^{NS}	-1.3209 ^{NS}	-0.1671	0.2165
7	Experience in cotton cultivation	-0.0868 ^{NS}	1.6167 ^{NS}	0.1895	0.1172
8	Annual income	0.3928**	0.4984 ^{NS}	0.0248	0.0498
9	Social participation	0.0515 ^{NS}	-0.8995 ^{NS}	-0.2926	0.3253
10	IPM equipment possession	0.4979**	0.4227 ^{NS}	0.0791	0.1871
11	Extension agency contact	0.5387**	0.4135 ^{NS}	0.0487	0.1178
12	Mass media exposure	0.5691**	1.6494 ^{NS}	0.2444	0.1482
13	Risk orientation	0.3290**	-0.7066 ^{NS}	-0.0899	0.1271
14	Economic motivation	0.3284**	1.6694 ^{NS}	0.1939	0.1161
15	Innovativeness	0.5686**	3.6497**	3.0458	0.8345
16	Attributes of IPM	0.0431 ^{NS}	-0.8712 ^{NS}	-0.4496	0.5161

** - Significant at 1 % level

* - Significant at 5 % level

R² = 0.5661

F value = 6.7693**

NS – Non-significant

CONCLUSION

Cotton is a very important commercial crop and is of vitally important both in the agricultural as well as industrial economy in a country. The farmers how to do knowledge was increased by attended IPM-FFS training conducted by state department of agriculture in the survey conducted area. It may be concluded from the study that when all the other variables are kept at constant level, a unit increase in educational status, mass media exposure and innovativeness increase in how to do knowledge among IPM oriented cotton growers under rainfed condition.

REFERENCES

1. Anonymous, (2001). Technology Mission on Cotton – Mini Mission II. Ministry of Agrl., GOI, CICR, Nagpur, 13 p.
2. Arulmurugan, S. (2000). Knowledge and Adoption of Post Harvest Technologies of Turmeric Growers. Unpub. M.Sc (Agri) Thesis. TNAU, Coimbatore.
3. Jayalakshmi, M. (2000). Existence of Knowledge and Technological gap between Systems nad the Contributing Factors – An Analysis. Unpub. M.Sc (Agri) Thesis. TNAU, Coimbatore.
4. Pretty, J., I. Guijit, I. Sconnes and J.Thompson (1992). “Regenerating Agriculture: The Agro-ecology of Low-External Input and Communication Development”. In: J: Holmberg, (eds.) Policies for a small IIED, London.
5. Rogers, E.M. (1983). Diffusion of Innovations, New York, The Free Press, 453p.
6. Venilla, M.A. (1998). Adoption of Innovations : Generated by Research System – An Analysis. Unpub. M.Sc (Agri) Thesis. TNAU, Coimbatore.

CITATION OF THIS ARTICLE

B. Sudhakar and K.A. Ponnusamy. Relationship and Influence Of Independent Variables On How To Do Knowledge Under Rainfed Condition-A Critical Analysis. Bull. Env. Pharmacol. Life Sci., Vol 8 [5] April 2019: 31-33