



Determination the Toxicity of Aflatoxin and T-2 Toxin Induced Stress on Immune Response System with feeding Activated Charcoal in broiler chicks

Muatsm Al-Masad¹ Karkaz. M. Thalij²

1. Department of Animal Production, Faculty of Agriculture, Jerash University, Jordan.

2. Department of Food Science and nutrition, Faculty of Agriculture, Jerash University, Jordan.

ABSTRACT

The activated charcoal (AC) 0.4% of feed were used with goal of providing further evidence as a factor for alleviation of toxicity and induced stress of aflatoxins (AF) and T-2 toxin on immune response system was examined in 160 male broiler chicks fed from 1 to 21 day of age. When adding 2.5 mg (AF) or 4 mg/kg T-2 toxin alone or in combination to the diet. At 21 day of age, 5 birds/pen were bled via cardiac puncture. Serum concentration of total protein, albumin, glucose, uric acid, triglycerides, cholesterol and alanine amino transferase (ALT), aspartate amino transferase (AST), lactate dehydrogenase (LDH) alkaline phosphatase (AP), and gamma glutamyl transferase (GGT), were determined. The other 5 birds/pen were intravenously injected with 4.0×10^6 Escherichia coli on day 21, then blood samples were collected at 60, 120 and 180 min post injection. Liver and spleen were collected after 180 min and colony forming units were determined. Results of the experiment indicated that AF or T-2 toxin alone or in combination caused a significant ($p < 0.05$) decrease in total protein, albumin, glucose, uric acid, triglycerides and cholesterol. The presence of these toxins alone or in combination, also resulted in significant decrease in enzyme activities of ALT, AST, LDH and AP except GGT was decreased in chicks fed AF or T-2 toxin only. There was significant increase in bacterial colonies counts in blood, liver and spleen compared with the control chicks group. The addition of activated charcoal at a concentration of 0.4% to the feed in this experiment was responsible for reduction of the toxic effects on all parameters when AF or T-2 toxin present individually or in combination in the diets.

Keywords: Aflatoxin, T-2 Toxin, Immune Response, Charcoal, broiler.

Received 22.07.2016

Revised 10.08.2016

Accepted 21.08.2016

INTRODUCTION

Mycotoxins are structurally diverse secondary metabolites of fungi that grow on a variety of feed and foods that consumed by animals and human, respectively. Consumption of lesser amounts of fungal toxins may result in impaired immunity and decreased resistance to infection disease, indeed, it has long been recognized by veterinary clinicians that marked immunosuppression is observed in livestock ingesting mycotoxins at levels below those that cause overt toxicity [1]. The aflatoxins (AF) are a group of closely related toxic metabolites produced in feedstuffs by *Aspergillus flavus* and *A. parasiticus*, plus related species *A. nomius* [2]. They commonly occur as natural contaminants of poultry feeds [3]. One of the myriad effects of mycotoxins is the ability to impair the immune system in fowl. Aflatoxin is the best-known mycotoxin for its ability to impair reticuloendothelial activity [3] (Bintvihok and Kositcharoenkul, 2007), primary immune response [4], complementary system [3], phagocytic activity of leukocytes and alveolar macrophages (Yunus et al, 2002; Muller et al, 2005). The inhibitory effect of aflatoxin on antibody production has also been demonstrated in chicks [4] and turkeys [5]. The T-2 toxin is a naturally occurring mycotoxin produced by several species of fungi in the genus *Fusarium* [6]. Those are found in many cereals, feeds, and vegetables. The T-2 toxin causes reduction in feed consumption and weight gain and severe lesions in chickens [7, 8], altered feathering [9], and a coagulopathy [7]. Both AF and T-2 toxin are important to the poultry industry due to their toxicity and occurrence in feeds, the toxicity of individual mycotoxins can be enhanced when they occur as co-contaminants of feeds [7]. Practical methods for detoxifying mycotoxins containing feeds on a large commercial scale and in a cost-effective manner are not currently available. A new approach to this problem has been the addition of non-nutritive sorptive materials to the diet known as sorbents in order to reduce the absorption of mycotoxins from the

gastrointestinal tract. In recent years, the last adsorbent was reported that was effective in reducing toxic of AF and T-2 toxin in broiler chicks [10]. Therefore, the objective of this study was to evaluate the ability of AC when included in a diet to protect AF and T-2 toxin in response immune system in broiler chicks.

MATERIALS AND METHODS

In this experiment 160 day-old male broiler chicks (Lohman) were housed in heated starter batteries under continuous fluorescent lighting. Chicks were fed a corn soybean meal based starter diet obtained from a commercial mill; it contained 22% CP and 2950 ME/kg. The diets were formulated without added antibiotics, coccidiosis or growth promoter. The chicks were randomly assigned to the following treatment groups: **1)** Control with 0 mg AF, 0 mg T-2, 0% AC; **2)** 0.04% AC; **3)** 2.5 mg AF/kg of diet; **4)** 2.5 mg AF/kg of diet plus 0.04% AC; **5)** 4 mg T-2/kg of diet; **6)** 4 mg T-2/kg of diet plus 0.04% AC; **7)** 2.5 mg AF plus 4 mg T-2/kg of diet; **8)** 2.5 mg AF plus 4 mg T-2/kg of diet plus 0.04% AC. Each treatment consisted of two replicates of ten chicks per replicate. Aflatoxin was prepared through inoculation of rice by *Aspergillus parasiticus* NRRL 2999 described by (Shotwell et al, 1966), as modified by (Nabney and Nisbitt, 1965). Fermented rice was autoclaved and ground and the AF content measured by Spectrophotometric analysis (Nabney and Nisbitt, 1965, Wiseman et al., 1967). The total AF content in the rice powder, 80% was AFB1, 14% was AFG1, 5% was AFB2 and 1% was AFG2. The T-2 toxin was prepared through inoculation of cracked corn by *Fusarium sporotrichioides* (Laboratory isolate) at 15 °C by method previously described by [10]. T-2 toxin was extracted and determined under the conditions described by [11] and incorporated into the diet by dissolving the toxin in 95% ethanol and then mixing the appropriate quantities with 1 kg of the diet. After drying, the dissolved toxin was mixed with the basal diet to produce the treatments containing T-2 toxin. The basal diet was analyzed for mycotoxins and was found to be below detection limits for AF, ochratoxin, zearalenone as established by the methods described by [12]. Feed and water were available for *ad libitum* consumption. At 21 days of age, five birds per pen were bled via cardiac puncture. The serum was collected using corvac serum separator tubes. Serum concentration of total protein, albumin, glucose, uric acid, triglycerides, cholesterol, alanine aminotransferase (ALT), aspartate aminotransferase (AST), Lactate dehydrogenase (LDH), γ -glutamyl transferase (GGT) and alkaline phosphatase (AP) were determined using a clinical chemistry analyzer according to the manufacturer's recommended procedure (obtained from Rondex Laboratories Ltd.). The other five birds were intravenously injected with 4.0×10^6 *E. coli*. The amount of *E. coli* used was based on pilot experiments, for this study, a dose was chosen that showed complete elimination of bacteria from circulation but did not cause severe hemodynamic changes influencing clearance function. One milliliter blood was collected from the jugular vein into heparinized tubes at 60, 120 and 180 min post injection. Prior to *E. coli* inoculation, blood samples from each bird were collected and checked for the presence of *E. coli* in order to insure no preexisting infection. At 180 min post injection, poulters were killed, and tissue samples of liver and spleen were taken for quantitative bacterial determination. Whole blood samples were serially diluted with sterile PBS and 100 μ l of each dilution at each time point was plated into MacConkey agar plates. Approximately one gram of tissue sample was homogenized with 5 ml of sterile PBS. Serial dilution of tissue suspension (100 μ l) was plated onto MacConkey agar plates. Plates were then incubated at 37°C for 18 h, and number of colonies on the plates were determined and expressed as *E. coli* colony forming units. Each sample run in duplicate plates. The final bacterial concentrations were calculated as the number of colonies per milliliters of blood and as colonies per gram of harvested tissue. The colonies grown on the plates were confirmed to be *E. coli* colonies by previously reported methods [13].

Data were analyzed by the ANOVA analysis, using the general linear model of the Statistically Analysis System [14]. Significant treatment differences were evaluated using Duncan's multiple-range test [15]. All statements of significance are based on the 0.5 level of probability.

RESULTS

The effects of dietary 2.5 mg AF or 4 mg T-2 toxin/kg of diet singly or in combination with or without 0.4% AC on serum biochemical values are summarized in table 1. When compared with the controls, serum total protein, albumin, glucose, uric acid, triglycerides, and cholesterol was significantly ($p < 0.05$) decreased in chicks fed diets containing AF or T-2 toxin singly or in combination. The addition of AC alone was not differ significantly when compared with the control group. Activated charcoal when added to the diet containing T-2 alone cause no change in these parameters except Albumin and glucose concentration that were ameliorative the negative effect of toxin. Also the negative effects which obtained in serum biochemical parameters caused when dietary AF was ameliorative by the addition of 0.5% AC to the chicks diets. The impact of adding the AC to chicks fed did not influence in complete protection from the

two toxins negative effects. But they were became intermediately effects between the toxicants treatment and control group.

Table 1. Individual and combined Effects of AF and T-2 toxin with or without AC' on blood biochemistry in broiler chicks.

AF	T-2	AC	TP*	Alb	Glu	UA	TG	Cho
mg/kg	%	gm/100 ml			mg/100 ml			
0	0	0	3.14 \pm 0.01	1.19 \pm 0.01	269 \pm 3.78	6.17 \pm 0.1	102 \pm 1.5	174 \pm 1.5
0	0	0.4	3.12 \pm 0.01	1.13 \pm 0.02	270 \pm 3.98	6.19 \pm 0.01	103 \pm 1.2	176 \pm 1.1
2.5	0	0	2.40 \pm 0.01	0.79 \pm 0.01	249 \pm 3.73	5.05 \pm 0.08	69 \pm 0.83	120 \pm 1.33
2.5	0	0.4	2.78 \pm 0.01	0.91 \pm 0.01	259 \pm 2.95	5.18 \pm 0.08	70 \pm 1.5	153 \pm 1.0
0	4	0	3.00 \pm 0.01	1.00 \pm 0.18	251 \pm 4.73	5.03 \pm 0.1	93 \pm 1.8	159 \pm 1.84
0	4	0.4	2.99 \pm 0.01	0.97 \pm 0.01	259 \pm 2.87	5.04 \pm 0.07	96 \pm 1.8	157 \pm 1.79
2.5	4	0	2.01 \pm 0.02	0.69 \pm 0.01	240 \pm 2.81	4.16 \pm 0.08	58 \pm 0.79	118 \pm 1.04
2.5	4	0.4	2.06 \pm 0.01	0.76 \pm 0.01	247 \pm 4.10	4.13 \pm 0.07	70 \pm 0.85	126 \pm 1.20

a-d Values within columns with no common superscript differ significantly at 0.05.

*AC=Activated charcoal. *TP=Total protein, Alb=Albumin, Glu=Glucose, UA=Uric acid, TG=Triglycerides, Cho=Cholesterols.

Data presented in table 2. Show the effects of dietary treatments on serum enzyme activities. The activity of alanine aminotransferase (ALT), aspartate aminotransferase (AST) lactate dehydrogenase(LDH), and alkaline phosphatase (AP), was significantly ($p < 0.05$) decreased, and the activity of γ -glutamyl transferase was increased in chicks fed diets containing AF or T-2 toxin individually or in combination, when compared with the control group. Adding AC to the diet containing AF alone or in combination with T-2 was acting in ameliorate the effects of the toxins on the enzyme activities. Whereas the added of AC to the diet containing T-2 alone did not effects on the activity of ALT enzyme.

Table 2. Individual and combined Effects of AF and T-2 toxin with or without AC' on serum Enzyme activities in broiler chicks.

AF	T-2	AC	ALT*	AST	LDH	GGT	AP
mg/kg	%	IU/ml					
0	0	0	20.5 \pm 0.33	145 \pm 1.42	444 \pm 5.8	10.4 \pm 0.20	5.99 \pm 0.01
0	0	0.4	20.4 \pm 0.17	146 \pm 1.40	443 \pm 4.8	10.4 \pm 0.14	6.03 \pm 0.01
2.5	0	0	10.5 \pm 0.16	87 \pm 1.46	357 \pm 4.1	13.4 \pm 0.13	4.11 \pm 0.01
2.5	0	0.4	14.0 \pm 0.20	121 \pm 0.92	392 \pm 4.0	11.6 \pm 0.23	4.85 \pm 0.01
0	4	0	18.1 \pm 0.10	134 \pm 0.67	320 \pm 3.1	13.0 \pm 0.11	3.91 \pm 0.01
0	4	0.4	17.9 \pm 0.10	137 \pm 0.68	384 \pm 5.0	11.0 \pm 0.11	3.85 \pm 0.01
2.5	4	0	9.1 \pm 0.10	83 \pm 1.46	320 \pm 3.95	11.5 \pm 0.20	3.79 \pm 0.01
2.5	4	0.4	10.3 \pm 0.05	85 \pm 0.74	302 \pm 5.40	10.6 \pm 0.15	3.78 \pm 0.01

a-d Values within columns with no common superscript differ significantly at 0.05.

*AC=Activated charcoal. *ALT = Alanine amino transferase. GGT = γ -glutamyl transferase. AST = Aspartate amino transferase. AP = Alkaline phosphatase. LDH = Lactate dehydrogenase.

Table 3. Individual and combined Effects of AF and T-2 toxin with or without AC' on *Escherichia coli* clearance from blood and tissue in broiler chicks.

AF	T-2	AC	Time post injection (log10cfu/ml blood)			(log10cfu/gm tissue after 180 min post injection)	
mg/kg	%		60 min	120 min	180 min	Liver	Spleen
0	0	0	2.80 \pm 0.03	2.27 \pm 0.01	1.02 \pm 0.04	4.55 \pm 0.04	4.88 \pm 0.03
0	0	0.4	2.81 \pm 0.01	2.29 \pm 0.01	1.02 \pm 0.01	4.56 \pm 0.01	4.90 \pm 0.01
2.5	0	0	3.28 \pm 0.01	3.26 \pm 0.01	3.18 \pm 0.01	5.02 \pm 0.01	6.02 \pm 0.01
2.5	0	0.4	3.07 \pm 0.01	3.11 \pm 0.09	2.88 \pm 0.01	4.80 \pm 0.01	5.85 \pm 0.01
0	4	0	2.90 \pm 0.01	2.50 \pm 0.01	1.40 \pm 0.01	4.59 \pm 0.01	5.14 \pm 0.01
0	4	0.4	2.90 \pm 0.01	2.48 \pm 0.01	1.39 \pm 0.01	4.57 \pm 0.01	5.12 \pm 0.01
2.5	4	0	3.40 \pm 0.01	3.37 \pm 0.01	3.29 \pm 0.01	5.02 \pm 0.01	6.08 \pm 0.01
2.5	4	0.4	3.28 \pm 0.01	3.24 \pm 0.01	3.19 \pm 0.01	4.89 \pm 0.01	5.90 \pm 0.01

a-d Values within columns with no common superscript differ significantly at 0.05.

•AC=Activated charcoal.

In comparison with the results in the control, the delayed systemic bacterial clearance in chicks fed 2.5 AF and 4 mg T-2 toxin/kg diet singly or in combination were associated with significantly higher numbers of viable bacteria in blood system and organs (Table 3). The accounts of *E. coli* colonies in blood system, liver, and spleen was associated with significantly higher accounts in chicks fed AF or T-2 toxin individually or in combination. The addition of AC did not effecting in complete protection as evidenced by all values which determined that were appear intermediate between those of control and AF and T-2 toxin singly or in combination.

DISCUSSION

The aflatoxins and T-2 toxin are important mycotoxins due to their toxicity and occurrence in feedstuffs that are used in poultry and livestock diets. In the present study, the toxic effects of AF or T-2 toxin singly or in combination were expressed as decreased concentration of serum total protein, albumin, and activity of alkaline phosphatase are most likely associated with inhibition of protein synthesis by AF through conversion to a 2,3 epoxide binding to DNA and inhibiting RNA synthesis [16, 17]. T-2 toxin also acted in inhibits the protein synthesis through the inactivation and termination process, possibly through its binding to ribosome's [18]. Therefore, aflatoxin primarily effects on protein synthesis during transcription, and T-2 toxin effects on protein synthesis during translation process, which may partially account for their synergistic toxicity. The reduction in concentration of serum cholesterol is most likely due to inhibition of cholesterol biosynthesis and the reduction in concentration of serum triglycerides might be associated with impaired lipid transport which causes from AF or T-2 toxin, as suggested by [19, 20]. The γ -glutamyl transferase activity is a sensitive indicator of liver disease, furthermore with other enzyme checked in this study, whether the disorder involves liver inflammation, a space occupying lesion, or obstruction of the biliary tract.

Reduction in systemic bacterial clearance observed in the current study may also be a result of dysfunction of the mononuclear phagocytic system [23, 24]. In addition to the delayed bacterial clearance in the blood, higher accounts of bacterial colonies in the tissue cultures of chicks fed AF or T-2 toxin singly or in combination also suggest that the ability to engulf bacteria is impaired. It is speculated that macrophages in mycotoxins fed chicks were unable to digest bacteria within macrophages, leading to higher accounts of bacteria present in the tissues. As a result, systematic bacterial clearance was delayed. The diminished systemic bacterial clearance was associated with increased of bacterial colonies accounts in tissues, indicating that the phagocytic and lytic capacity of the mononuclear phagocytic system was compromised [21,22].

The results indicate that the AC tested was marginally effective in reducing some of the toxic signs of chronic toxicosis in growing broilers fed AF, but of little benefit when T-2 toxin was fed. Any protective effect probably involves the sequestration of the toxic molecules in the gastrointestinal tract and chemisorption to the charcoal.

REFERENCES

1. Oswald, I.P., (2007). Effects immunosuppresses des mycotoxins chez le porc. Journées Recherché Porcine, 39, 419-426.
2. Binder, E.M., L.M. Tan, L.J. Chin, J. Handle and J. Richard, (2007). Worldwide occurrence of mycotoxins in commodities feed and feed ingredients. Anim. Feed Sci. Technol., 137: 265-282.
3. Bintvihok, A., Kositcharoenkul, S., (2006). Effect of dietary calcium propionate on performance, hepatic enzyme activities and aflatoxin residues in broilers fed a diet containing low levels of aflatoxin B1. *Toxicon*, 47, 41-46.
4. Celik, I., Oguz, H., Demet, O., Donmez, H.H., Boydak, M., Sur, E., (2000). Efficacy of polyvinylpyrrolidone in reducing the immunotoxicity of aflatoxin in growing broilers. *British Poultry Science*, 41, 430-439.
5. García, J.C.D.R.; Ramos, C.M.; Pinton, P.; Elvira, S.M.; Oswald, I.P., (2007). Evaluation of the cytotoxicity of AFB1, FB1 and AFB1/FB1 in intestinal cell. *Revista Iberoamericana de Micologia*, 24, 136-141.
6. Diaz, D.E., 2005. The Mycotoxin Blue Book, 1st ed.; Nottingham University Press: Nottingham, UK, pp. 25-56.
7. Girish, CK, Devegowda, G., (2006). Efficacy of glucomannancontaining yeast product (Mycosorb®) and hydrated sodium calcium aluminosilicate in preventing the individual and combined toxicity of aflatoxin and T-2 toxin in commercial broilers. *Asian-Aust J Anim Sci.*, 19:877-83.
8. Kubena LF, Harvey RB, Huff WE, Corrier TD, Phillips TD, Rottinghaus GE., (1990). Efficacy of hydrated sodium calcium aluminosilicate to reduce the toxicity of aflatoxin and T-2 toxin. *Poult Sci*, 69: 1078-1086.
9. Raju, M.V.L.N. and G. Devegowda, (2000). Influence of esterified glucomannan on performance and organic morphology, serum biochemistry and hematology in broilers exposed to individual and combined mycotoxicosis (aflatoxin, ochratoxin and T-2 toxin). *British Poult. Sci.*, 41: 640-650.
10. Edrington TS, Kubena LF, Harvey RB, Rottinghaus GE., (1997). Influence of a super activated charcoal on the toxic effects of aflatoxins or T-2 toxin in growing broilers. *Poultry Sci.*, 76: 1205-1211.

11. Cullen D , (1982). Smalley EB , Caldwell RW. New process for T-2 toxin production. *Appl Environ Microbiol*;;44:371- 375.
12. Clement BA, Phillips TD., (1985). Advances in the detection and determination of mycotoxins via capillary GC/quadrupole mass spectrometry. *Toxicologist*;; 5(1):232.
13. Patten VH , Shin SJ, Cole J, Waston CW , Fales WH., (1995). Evaluation of a commercial automated system and soft-ware for identification of veterinary bacterial isolates. *J Vet Diagn Invest*;; 7: 506-508.
14. Statistical Analysis System SAS Users guide: Cary, NC. 27512 – 8000, (2001); U.S.A.
15. Duncan, DB. (1955). Multiple range and F; test. *Biometrics*;; 11:1-42.
16. Miazzo, R., Peralta, M.F. ; Magnoli, C.; Salvano, M.; Ferrero, S.; Chiacchiera, S.M.; Carvalho, E.C.Q.; Rosa, C.A.R. and Dalcero, A., (2005). Efficacy of sodium bentonite as a detoxifier of broiler feed contaminated with aflatoxin and fumonisin. *Poult. Sci.*, 84: 1-8.
17. Yu FL. (1981). Studies on the mechanism of aflatoxin B1 inhibition of rat liver nucleolar RNA synthesis. *J Biol Chem*;; 256:3292-3297.
18. Ueno Y., (1977). Mode of action of trichothecens. *Am Nutr Alim*;; 31:885-900.
19. Manafi M., M.; Umakantha, B. Ali, N and Swamy, H.D., (2012). Study of the Combination Effects of Aflatoxin and T-2 Toxin on Performance Parameters and Internal Organs of Commercial Broilers. *Global Veterinaria* 8 (4): 393-396.
20. Kubena LF, Smith EE, Gentles A, Harvey RB, Edrington TS, Phillips TD, Rottinghaus GE., (1994). Individual and combined toxicity of T-2 toxin and cyclopiazonic acid in broiler chicks. *Poult Sci*;; 73:1390-1397.
21. Edrington TS, Sarr AB, Kubena LF, Harvey RB, Phillips TD., (1996). Effects of hydrated sodium calcium aluminosilicate (HSCAS), acidic HSCAS and activated charcoal, on the metabolic profile and toxicity of aflatoxin B1 in turkey poults. *Toxicol. Lett*;; 89:115-122.
22. Miazzo, R.; Rosa, C.; de Queiroz, C.E.; Magnoli, C.; Chiacchiera, S.; Palacio, G.; Saenz, M.; Kikot, A.; Basaldella, E.; Dalcero, A., (2000). Efficacy of synthetic zeolite to reduce the toxicity of aflatoxin in broiler chicks. *Poult. Sci.*, 79, 1–6.
23. Fairchild, A.S.; Croom, J.; Grimes, J.L. and Hagler, W.M., 2008. Effect of ASTRA-BEN 20 on Broiler Chicks Exposed to Aflatoxin B or T2 Toxin, *International Journal of Poultry Science* 7 (12): 1147-1151.
24. Qureshi MA , Brake J, Hamilton PB, Hagler, Jr WM., Nesheim S., (1998). Dietary exposure of broiler breeders to aflatoxin results in immune dysfunction in progeny chicks. *Poult Sci*;; 77:812-819.

CITATION OF THIS ARTICLE

Muatsm Al-Masad and Karkaz. M. Thalij. Determination the Toxicity of Aflatoxin and T-2 Toxin Induced Stress on Immune Response System with feeding Activated Charcoal in broiler chicks. *Bull. Env. Pharmacol. Life Sci.*, Vol 5 [10] September 2016: 59-63