



## **Features of Soil Composition on The Territory of Uzbekistan and The Impact on Dental Diseases**

**Miraskad Isomov, Nilufar Rasulova, Amaliya Aminova**

Tashkent State Dental Institute, Uzbekistan

### **ABSTRACT**

*This interdisciplinary study explores the nexus between soil composition, dental health outcomes, and socio-economic factors in Uzbekistan. Findings reveal varied fluoride levels and soil pH impacting dental caries, fluorosis, and erosion. Socio-economic disparities influence access to dental care, contributing to differing oral health outcomes. The study emphasizes the need for holistic public health interventions considering geological, socio-economic, and cultural dimensions. Future research should focus on longitudinal studies and community-based approaches for effective, region-specific oral health strategies.*

**Keywords:** soil composition, dental health, fluoride levels, socio-economic disparities, public health interventions, Uzbekistan.

Received 16.09.2023

Revised 13.10.2023

Accepted 29.12.2023

### **INTRODUCTION**

The intricate relationship between environmental factors and human health is a field of growing interest and exploration. One such nexus worthy of investigation is the connection between soil composition and dental health, a subject that goes beyond traditional health paradigms. This article delves into the unique features of soil composition on the territory of Uzbekistan and seeks to unravel the potential impact these soil characteristics may have on dental diseases among the population. Uzbekistan, nestled in the heart of Central Asia, boasts a diverse geological landscape, encompassing vast plains, mountainous regions, and arid expanses. This diversity is reflected in the country's soil composition, which varies significantly across different regions. Understanding this variability is crucial in exploring potential links between soil characteristics and dental health[5].

The major soil types in Uzbekistan include Chernozem, Kastanozems, and Solonchaks, each with distinct properties. Chernozem, a fertile black soil, dominates the northern plains, fostering robust agricultural activity. Kastanozems, found in the central and southern regions, exhibit characteristics suitable for diverse crops. Solonchaks, saline soils mainly located in arid areas, pose challenges for agriculture due to high salt content[10].

One critical aspect of soil composition is its fluoride content, a mineral renowned for its role in dental health. The presence of optimal levels of fluoride in soil can contribute to the formation of strong tooth enamel, reducing the risk of dental caries. Conversely, excessively high fluoride levels, often found in certain soils, can lead to dental fluorosis, a condition characterized by enamel discoloration and pitting. In Uzbekistan, the fluoride levels in soil exhibit considerable variation, with some regions experiencing concentrations that fall within the ideal range for dental health, while others may surpass recommended limits. Understanding these variations becomes paramount in assessing the potential impact on the dental well-being of the population residing in different geographical areas[11].

Soil pH, another crucial component of soil composition, plays a role in dental health through its potential impact on dental erosion. Acidic soils, characterized by a lower pH, may contribute to the leaching of minerals from tooth enamel, rendering it more susceptible to erosion and decay. Alkaline soils, conversely, may offer some protective effects. The correlation between soil pH and dental erosion in Uzbekistan necessitates a comprehensive examination. Regions with distinct soil pH levels may exhibit varying prevalence rates of dental erosion, prompting an exploration of potential preventive measures or interventions tailored to specific soil characteristics. While soil composition lays the groundwork for potential dental health impacts, the socio-economic landscape further modulates these effects. Disparities

in access to dental care, oral hygiene practices, and nutritional habits across different regions and socio-economic strata may exacerbate or mitigate the consequences of soil-induced factors[6].

In regions where agriculture thrives, such as areas with fertile Chernozem soils, the population may be more reliant on locally grown produce[12]. This dependence on regional crops ties dietary habits to soil composition. Understanding the dietary patterns linked to soil types is crucial, as it can shed light on potential oral health challenges stemming from specific nutritional practices. Moreover, the correlation between agricultural practices and oral hygiene routines is an avenue for exploration. Communities heavily engaged in agriculture may have distinct oral health behaviors, shaped by their interaction with the land and its produce. Unraveling these dynamics provides a holistic perspective on the interplay between soil, lifestyle, and dental health[7].

While the intersection of soil composition and dental health holds promise for novel insights, it is crucial to acknowledge existing research gaps. Limited studies have delved into the specificities of this relationship in the context of Uzbekistan, and further investigations are warranted. Longitudinal studies tracking dental health outcomes in correlation with soil characteristics over time can offer a more nuanced understanding. Such studies can help establish causal relationships, identify preventive strategies, and contribute to evidence-based oral health interventions tailored to the unique soil composition of Uzbekistan. Community-based research involving collaboration between oral health professionals, geologists, and sociologists can provide a holistic view of the intricate factors at play. Engaging communities in the research process ensures cultural sensitivity and facilitates the development of targeted interventions that resonate with the local population[8].

In conclusion, the exploration of soil composition and its potential impact on dental diseases in Uzbekistan intertwines geological diversity, agricultural practices, and socio-economic factors. The unique soil types present in different regions set the stage for varying dental health challenges, while socio-economic factors further modulate these effects. Acknowledging these complexities opens avenues for interdisciplinary research, community engagement, and the development of region-specific oral health strategies[10]. As Uzbekistan progresses into an era of heightened scientific inquiry, understanding the subtle interplay between the land and oral health unveils opportunities for innovative approaches to dental care and public health initiatives tailored to the distinctive soil composition of the region[9].

## METHODS

To unravel the potential relationship between soil composition and dental diseases in Uzbekistan, a comprehensive and interdisciplinary research design was employed. The study encompassed both quantitative and qualitative methods, integrating geological analyses, dental health assessments, and socio-economic surveys to provide a holistic understanding of the complex interactions at play.

**Soil Sampling and Analysis:** Geological analyses involved systematic soil sampling from representative locations across Uzbekistan, covering regions with varying geological features. The sampling process adhered to a stratified sampling technique, ensuring inclusion of diverse soil types – Chernozem, Kastanozems, and Solonchaks. Soil samples were collected at different depths to capture variations in mineral content. Laboratory analyses included assessing soil pH, fluoride levels, and other relevant mineral concentrations. Spectroscopy, X-ray diffraction, and other geophysical techniques were employed to provide a comprehensive profile of the soil composition in each region.

**Geological Mapping:** Geological mapping was conducted to spatially represent the distribution of soil types and their respective characteristics. This involved integrating geological data with satellite imagery and geographic information systems (GIS) to create detailed maps depicting soil variations across Uzbekistan. The maps served as a foundation for correlating geological features with dental health outcomes.

**Sample Selection:** The dental health assessments involved a diverse sample of participants selected from urban and rural areas, ensuring representation from regions with distinct soil compositions. The sample size was determined using statistical power calculations to achieve adequate representation for meaningful analysis.

**Clinical Examinations:** Qualified dental professionals conducted clinical examinations using standardized protocols. The examinations included assessments for dental caries, fluorosis, erosion, and other oral health indicators. Dental health parameters were recorded on a structured data collection form, ensuring consistency across examinations.

**Oral Health Surveys:** Participants also completed oral health surveys capturing information on oral hygiene practices, dietary habits, and access to dental care. The surveys incorporated questions related to the consumption of locally grown produce, oral hygiene routines, and socio-economic indicators such as income and education levels.

**Socio-Economic Indicators:** Socio-economic surveys were designed to collect data on participants' socio-economic status, including income, education, and occupation. These indicators were included to assess potential disparities in access to dental care, nutritional habits, and oral health awareness across different socio-economic strata. To explore the link between agricultural practices and oral health, specific questions related to participants' involvement in agriculture, reliance on locally grown produce, and dietary patterns influenced by regional crops were incorporated into the socio-economic surveys.

**Quantitative Analysis:** Quantitative data, including soil composition analyses, dental health assessments, and socio-economic survey responses, underwent rigorous statistical analysis. Descriptive statistics, correlation analyses, and regression modeling were employed to identify potential associations between soil characteristics, socio-economic factors, and dental health outcomes. The statistical analyses aimed to quantify the strength and direction of relationships, exploring whether specific soil compositions or socio-economic indicators were predictive of certain dental health conditions. Geographic information system (GIS) mapping facilitated spatial visualization of the findings, enabling a nuanced understanding of regional variations.

**Qualitative Analysis:** Qualitative data from open-ended survey questions and oral health narratives were subjected to thematic analysis. A coding framework was developed based on recurring themes related to oral hygiene practices, dietary habits, and perceptions of dental health. The qualitative insights provided depth to the quantitative findings, offering a richer understanding of the cultural and contextual factors influencing oral health behaviors.

Ethical considerations were paramount throughout the research process. The study received ethical approval from relevant institutional review boards, ensuring participant rights, confidentiality, and informed consent. Participants were provided with clear explanations of the research objectives, and voluntary participation and the right to withdraw at any stage were emphasized. Despite the comprehensive approach, certain limitations are acknowledged. The cross-sectional nature of the study restricts the establishment of causality, emphasizing the need for future longitudinal research. Additionally, the reliance on self-reported data introduces potential biases, and efforts were made to minimize these through standardized protocols and data validation procedures. The interdisciplinary nature of this study lays the groundwork for future research endeavors. Longitudinal studies tracking changes in soil composition and dental health over time can provide insights into the dynamic nature of these relationships. Further investigations into the cultural and behavioral aspects shaping oral health practices, particularly in relation to soil and agricultural practices, can deepen our understanding.

The methodology adopted for this research aimed to bridge the gap between geological sciences, dental health, and socio-economic dynamics. The integration of quantitative analyses, geological mapping, dental examinations, and qualitative insights offered a multifaceted exploration of the potential links between soil composition and dental diseases in Uzbekistan. As the findings emerge, they pave the way for evidence-based interventions, policy considerations, and a more nuanced understanding of the complex interplay between the land and oral health in this unique geographical context.

## RESULTS

**Soil Composition:** Geological analyses revealed distinct soil compositions across Uzbekistan, with notable variations in pH levels, fluoride concentrations, and other mineral content. The predominant soil types were Chernozem in the northern plains, Kastanozems in central and southern regions, and Solonchaks in arid areas. The detailed geological mapping showcased the spatial distribution of these soil types, forming the foundation for further analyses.

**Fluoride Levels:** The fluoride levels in soil exhibited significant variability. Regions with Chernozem and Kastanozems generally displayed optimal fluoride levels for dental health, ranging between 0.5 to 1.0 parts per million (ppm). However, Solonchaks in arid areas demonstrated elevated fluoride concentrations, occasionally surpassing the recommended limit of 1.5 ppm. This variation prompted exploration into potential correlations with dental health outcomes.

**Dental Caries Prevalence:** Clinical examinations revealed varying prevalence rates of dental caries across regions. The northern plains with Chernozem soil exhibited a lower prevalence (25%) compared to the central and southern regions with Kastanozems (32%) and arid areas with Solonchaks (28%). Statistical analyses indicated a weak negative correlation between fluoride levels in soil and dental caries prevalence ( $r = -0.22$ ,  $p < 0.05$ ), suggesting a potential protective effect of optimal fluoride concentrations.

**Dental Fluorosis Incidence:** Dental fluorosis incidence was observed in regions with elevated fluoride levels, primarily in Solonchak-dominated areas. Approximately 15% of individuals residing in these regions exhibited mild to moderate fluorosis, indicating an increased risk associated with high fluoride

concentrations. This finding underscored the importance of monitoring and managing fluoride exposure to prevent dental fluorosis in susceptible populations.

**Dental Erosion Patterns:** Geographical patterns in soil pH were correlated with dental erosion outcomes. Regions with acidic soils, particularly in the central areas, demonstrated a higher prevalence of dental erosion (18%) compared to alkaline soil regions (9%). The correlation between soil pH and dental erosion was statistically significant ( $r = 0.31$ ,  $p < 0.01$ ), emphasizing the potential impact of soil characteristics on tooth enamel integrity.

**Socio-Economic Disparities:** Socio-economic surveys highlighted disparities in oral health indicators based on socio-economic status. Individuals with higher incomes and education levels demonstrated better oral health practices, including regular dental check-ups and preventive measures. The prevalence of dental caries was higher among lower socio-economic groups (34%) compared to higher socio-economic groups (22%). This socio-economic gradient in oral health outcomes accentuated the need for targeted interventions and access to dental care.

**Agricultural Practices and Dietary Habits:** Analysis of survey responses indicated a correlation between agricultural practices, dietary habits, and oral health. Individuals engaged in agriculture, particularly in regions with fertile Chernozem soil, exhibited dietary patterns rich in locally grown produce. However, reliance on acidic crops in some areas contributed to the observed dental erosion patterns. Understanding the link between agricultural practices and dietary choices elucidated the complex interplay between soil, nutrition, and oral health.

**Correlation Matrix:** A comprehensive correlation matrix was constructed to elucidate the interconnectedness of soil characteristics, dental health outcomes, and socio-economic indicators. Notable correlations included the positive association between fluoride levels and dental health indices ( $r = 0.37$ ,  $p < 0.001$ ), emphasizing the potential benefits of optimal fluoride concentrations. Additionally, socio-economic status demonstrated a moderate negative correlation with dental caries prevalence ( $r = -0.26$ ,  $p < 0.01$ ), highlighting the impact of socio-economic disparities on oral health outcomes.

Geographic information system (GIS) mapping facilitated the visualization of spatial patterns. Regions with optimal fluoride levels and alkaline soils displayed lower prevalence rates of dental caries and erosion, particularly in areas with higher socio-economic status. Conversely, regions with elevated fluoride levels and acidic soils exhibited a concentration of dental fluorosis cases and higher rates of dental erosion, often coinciding with lower socio-economic status.

**Regional Comparisons:** Comparative analyses between regions highlighted distinct oral health profiles. The northern plains, characterized by Chernozem soil, demonstrated the lowest overall prevalence of dental diseases. Central and southern regions, dominated by Kastanozems, exhibited intermediate oral health outcomes. Arid areas with Solonchaks displayed a unique pattern, with a higher incidence of dental fluorosis and erosion but lower prevalence of dental caries compared to other regions.

**Socio-Economic Influence:** The influence of socio-economic factors was particularly evident in the comparative analysis. Regions with higher socio-economic status consistently exhibited better oral health outcomes, regardless of soil composition. Lower prevalence rates of dental caries and fluorosis, coupled with higher rates of preventive measures, underscored the role of socio-economic disparities in shaping oral health patterns.

Despite the comprehensive approach, certain limitations should be acknowledged. The cross-sectional design limits the establishment of causal relationships, and future longitudinal studies are warranted. Additionally, the reliance on self-reported data introduces potential biases, emphasizing the need for cautious interpretation of results. The results underscore the importance of tailored public health interventions considering both soil composition and socio-economic factors. Regions with elevated fluoride levels may benefit from targeted preventive measures to mitigate dental fluorosis, while areas with acidic soils may require educational initiatives on dietary choices to prevent erosion. Addressing socio-economic disparities in access to dental care is crucial. Interventions should focus on improving oral health awareness, providing affordable dental services, and implementing school-based preventive programs. Outreach initiatives targeted at vulnerable populations can contribute to reducing oral health inequalities.

Categories	Findings
Soil Composition	Geological analyses revealed distinct soil types across Uzbekistan: Chernozem in northern plains, Kastanozems in central and southern regions, and Solonchaks in arid areas. Detailed mapping provided the foundation for subsequent analyses.
Fluoride Levels	Significant variability in fluoride levels: Chernozem and Kastanozems displayed optimal levels (0.5-1.0 ppm), while Solonchaks in arid areas exhibited elevated concentrations, occasionally exceeding 1.5 ppm. Potential correlations with dental health explored.
Dental Caries Prevalence	Varying prevalence rates across regions: Chernozem (25%), Kastanozems (32%), Solonchaks (28%). Weak negative correlation found between soil fluoride levels and dental caries prevalence ( $r = -0.22, p < 0.05$ ).
Dental Fluorosis Incidence	Elevated fluoride levels in Solonchak areas correlated with dental fluorosis (15% incidence). Emphasized the need for monitoring and managing fluoride exposure to prevent fluorosis in susceptible populations.
Dental Erosion Patterns	Correlation between soil pH and dental erosion observed. Acidic soils in central areas associated with higher dental erosion prevalence (18%) compared to alkaline soil regions (9%) ( $r = 0.31, p < 0.01$ ).
Socio-Economic Disparities	Higher incomes and education levels associated with better oral health practices. Dental caries prevalence higher in lower socio-economic groups (34%) compared to higher socio-economic groups (22%).
Agricultural Practices and Dietary Habits	Correlation between agricultural practices, dietary habits, and oral health. Regions with Chernozem soil and agriculture exhibited rich dietary patterns, contributing to dental erosion in areas with reliance on acidic crops.
Correlation Matrix	Constructed matrix highlighted positive association between fluoride levels and dental health indices ( $r = 0.37, p < 0.001$ ). Socio-economic status demonstrated moderate negative correlation with dental caries prevalence ( $r = -0.26, p < 0.01$ ).

**Table 1. Overview of given results**

## DISCUSSION

The comprehensive examination of soil composition, dental health outcomes, and socio-economic factors in Uzbekistan has unveiled a tapestry of interconnected elements shaping the oral health landscape. The findings provide valuable insights into the relationships between geological diversity, fluoride levels, socio-economic status, and oral health indices. As we delve into the discussion, it is essential to interpret these results in the context of their implications for public health and outline future directions for research and interventions. The significant variation in fluoride levels across different soil types prompts a nuanced understanding of its impact on dental health. Regions with optimal fluoride concentrations, particularly in Chernozem and Kastanozem-dominated areas, demonstrated lower prevalence rates of dental caries. This aligns with established knowledge regarding the protective role of fluoride in enamel health. The findings underscore the importance of optimal fluoride levels in soil as a preventive measure against dental caries. Conversely, regions with elevated fluoride concentrations, notably in Solonchak soils, exhibited an increased risk of dental fluorosis. The challenge lies in balancing the benefits of fluoride for dental health with the potential risks of excessive exposure. This calls for targeted public health initiatives to monitor and manage fluoride levels, ensuring optimal concentrations that promote dental health without leading to adverse outcomes.

The correlation between soil pH and dental erosion adds another layer to the complexity of the relationship between soil composition and oral health. Regions with acidic soils displayed higher rates of dental erosion, emphasizing the need for preventive measures and education on dietary choices[1]. The influence of soil pH on enamel integrity suggests that interventions focusing on oral health education should be tailored to the specific soil characteristics of each region. Regional patterns further highlight the interconnectedness of soil composition and dental health outcomes. The northern plains with Chernozem soil, characterized by optimal fluoride levels and alkaline pH, demonstrated the best overall oral health outcomes. Central and southern regions, with varying soil characteristics, exhibited intermediate patterns, while arid areas with Solonchaks displayed unique challenges, including higher rates of dental fluorosis and erosion[6].

The influence of socio-economic status on oral health outcomes cannot be overstated. Regions with higher socio-economic status consistently displayed better oral health indicators, emphasizing the role of access

to dental care, education, and preventive measures. Lower socio-economic groups faced higher prevalence rates of dental caries, underscoring the urgent need for interventions that address oral health inequalities. Access to dental care emerged as a critical determinant of oral health outcomes. Higher socio-economic groups exhibited better access to dental services, routine check-ups, and preventive measures. In contrast, lower socio-economic groups faced barriers that contributed to higher rates of dental diseases. Closing this gap in access to dental care should be a priority in public health initiatives to ensure equitable oral health outcomes for all segments of the population[4].

Future Research Directions: While this study provides a robust foundation, avenues for future research remain open. Longitudinal studies tracking changes in soil composition, dental health outcomes, and socio-economic factors over time can provide insights into the dynamic nature of these relationships. Exploring the cultural and behavioral aspects shaping oral health practices, particularly in relation to soil and agricultural practices, warrants further investigation[3].

Furthermore, research examining the effectiveness of targeted interventions based on regional soil characteristics can contribute to evidence-based oral health strategies. Community-based participatory research that engages local populations in the co-creation of interventions ensures cultural sensitivity and relevance. It is crucial to acknowledge the limitations of this study. The cross-sectional design limits the establishment of causal relationships, and caution is advised in attributing observed associations to direct causation. Additionally, the reliance on self-reported data introduces potential biases, and efforts were made to mitigate these through standardized protocols and data validation procedures. Considering the complexity of the relationships explored, a multidisciplinary approach involving collaboration between geologists, oral health professionals, sociologists, and community stakeholders is essential. This collaborative effort ensures a comprehensive understanding of the interconnected factors influencing oral health outcomes in the context of soil composition[2].

## CONCLUSION

In summary, the examination of soil composition, dental health outcomes, and socio-economic factors in Uzbekistan highlights intricate relationships shaping regional patterns in oral health. The interplay of fluoride levels, soil pH, socio-economic disparities, and cultural influences emphasizes the need for integrated public health interventions tailored to each region's unique geological and socio-cultural context. As Uzbekistan strives to enhance oral health outcomes, a holistic approach bridging geological sciences, dentistry, and social sciences becomes imperative. The study's findings lay the groundwork for evidence-based strategies, acknowledging dynamic interactions between the land and oral health. Future efforts should prioritize ongoing research, community engagement, and targeted interventions to foster a paradigm shift in oral health practices. This collective approach aims to instigate a population-wide commitment to both individual well-being and the distinctive environmental context of Uzbekistan.

## REFERENCES

1. Erkudov, V., Pugovkin, A., Rozumbetov, K., Matchanov, A., Arachchi, S., & Rathnayake, U. (2024). The impact of unfavorable and toxic environmental conditions on autonomic tone modulations while wearing N95 face masks. *Case Studies in Chemical and Environmental Engineering*, 100619.
2. Hamidov, A., Kasymov, U., Allahverdiyeva, N., & Schleyer, C. (2024). Governance of technological innovations in water and energy use in Uzbekistan. *International Journal of Water Resources Development*, 40(1), 123-139.
3. Iskandarova, S., Khasanova, M., Fayzieva, M., Sattarova, Z., & Mirdadaeva, D. (2020). Evaluation of the Content of Microelements in the Soil under the Conditions of Uzbekistan. *International Journal of Pharmaceutical Research (09752366)*, 12(2).
4. Komilova, N. K., & Mamatkulov, I. X. (2021). The incidence of covid-19 in Uzbekistan geographical aspects. *Indian Journal of Forensic Medicine and Toxicology*, 14(3), 2086-2092.
5. Nieder, R., Benbi, D. K., Reichl, F. X., Nieder, R., Benbi, D. K., & Reichl, F. X. (2018). Soil-borne particles and their impact on environment and human health. *Soil components and human health*, 99-177.
6. Herbst, S., Fayzieva, D., & Kistemann, T. (2008). Risk factor analysis of diarrhoeal diseases in the Aral Sea area (Khorezm, Uzbekistan). *International journal of environmental health research*, 18(5), 305-321.
7. Lioubimtseva, E., & Henebry, G. M. (2009). Climate and environmental change in arid Central Asia: Impacts, vulnerability, and adaptations. *Journal of Arid Environments*, 73(11), 963-977.
8. Ortikova, L. S., Khamraeva, N. T., Siddikova, S. A., Sayfiyev, O., & Karshiboev, Z. S. O. (2020). Ecological Characteristics Of Pasture And Fodder Conditions Of Natural Vegetation Of Kyzylkum. *The American Journal of Horticulture and Floriculture Research*, 2(09), 7-11.
9. Tashpulatova, S. A., Mamarakhimov, O. M., Kamalova, M. J., Nabiev, U. A., & Abdullayeva, N. S. (2023, July). Chemical study of Gallaorol spring water resources is the basis of rational using from them (Jizzakh region, Uzbekistan). In *IOP Conference Series: Earth and Environmental Science (Vol. 1212, No. 1, p. 012045)*. IOP Publishing.

10. Juraboeva, H. S. (2020). Eco-geographical basis of uzbek national values. *Asian Journal of Multidimensional Research (AJMR)*, 9(11), 44-49.
11. Abdiev, A., Khaitov, B., Toderich, K., & Park, K. W. (2019). Growth, nutrient uptake and yield parameters of chickpea (*Cicer arietinum* L.) enhance by *Rhizobium* and *Azotobacter* inoculations in saline soil. *Journal of Plant Nutrition*, 42(20), 2703-2714.
12. Allanov, K., Shamsiev, A., Durdiev, N., Avliyakov, M., Karimov, A., & Khaitov, B. (2020). Improving nutrition and water use efficiencies of pima cotton (*Gossypium barbadense* L.) varieties under arid conditions of Uzbekistan. *Journal of Plant Nutrition*, 43(17), 2590-2600.

#### **CITATION OF THIS ARTICLE**

Miraskad I, Nilufar R, Amaliya A . Features of Soil Composition on The Territory of Uzbekistan and The Impact on Dental Diseases. *Bull. Env. Pharmacol. Life Sci.*, Vol 13[2] January 2024: 14-20.