



Scrotal Pathology Evaluation Using High-Resolution Sonography Imaging

Ashish Kumar Gupta, Srikanth Sanjeev Shetty, G. Jayalakshmi

Department of Microbiology, Sri Lakshmi Narayana Institute of Medical Sciences Affiliated to Bharath Institute of Higher Education and Research, Chennai, Tamil Nadu, India.

CORRESPONDING: deanuniversityaffairsslims@gmail.com, jayalakshmi.2k15@gmail.com

ABSTRACT

The primary goal of this study is to learn more about scrotal pathology and research it. It has been discovered that there has been significant progress in terms of both diagnosis and treatment. In this regard, it is essential to determine what factors influence scrotal diseases and to identify boxes, tables, and photos that might aid clinical analysis. This study aims to see how real-time grayscale sonography works in detecting scrotal lesions and what imaging properties it has. Furthermore, we try to distinguish between benign and malignant scrotal masses and correlate sonographic results with guided FNAC and/or biopsy wherever possible.

Keywords: Pathology; Scrotal symptoms; Sonography; Imaging techniques.

Received 11.09.2022

Revised 21.10.2022

Accepted 05.11.2022

INTRODUCTION

In clinical practice, scrotal edoema and discomfort are very common [1]. Although a patient's history and physical examination may be sufficient to make a diagnosis in certain cases, many patients require further tests to get a comprehensive picture of their symptoms [2, 3]. Clinical examinations are frequently deceptive or non-specific. The presence of discomfort and edoema of the scrotal contents might sometimes prevent a thorough examination [4]. Patients with sudden onset of scrotal discomfort might be difficult to diagnose. Early exploration has traditionally been favoured [5]. However, because it is believed that over 80% of acute scrotal processes are inflammatory in origin, this may result in a slew of unnecessary procedures.

Research published in 1974 was the first to employ B-mode ultrasonography to assess the testes and scrotum [6]. Ultrasonography has become an invaluable diagnostic technique for evaluating scrotal disease due to dramatic sonographic technology and apparatus developments over the previous two decades [7]. In the identification, characterization, and localization of intrascrotal anomalies, high-resolution real-time ultrasonography has proven a high degree of accuracy and sensitivity [8]. Noninvasive imaging and evaluation of tiny arteries in superficial organs is now achievable because too high-resolution real-time ultrasonography. The testis is the superficial organ where CDFI has had the most impact too far. It has reduced the number of clinical diagnostic options evaluated in the differential diagnosis, reducing the necessity for scrotal surgery and assisting in making precise diagnoses in situations of acute scrotal discomfort [9].

Because computed tomography uses ionizing radiation, it is not advised for gonadal evaluation [10]. CT's use in scrotal pathology is limited to staging testicular cancers by determining the precise location of retroperitoneal lymph nodes [11]. The normal scrotum and its contents may be seen in fine anatomic detail using a Magnetic Resonance Imaging (MRI) session [12]. Its large field of view allows it to check both the right and left hemi scrotal contents and the inguinal areas simultaneously, which is a significant benefit over ultrasonography. The testes, epididymis, and spermatic cord can all be distinguished thanks to the great contrast and spatial resolution [13]. The costs of MRI, as well as the lengthier examination duration and picture deterioration caused by patient mobility, are drawbacks.

In the diagnosis of testicular torsion, testicular scintigraphy has good sensitivity and specificity [14]. Its low resolution, lack of structural information, high cost, and use of ionizing radiation, however, restrict it. The advantages of both testicular scintigraphy and grey scale sonography are combined in colour Doppler flow

imaging [15]. The goal of this study was to assess the use and accuracy of high-resolution sonography and use this multifaceted data to lessen diagnostic challenges in these instances.

MATERIAL AND METHODS

The research was carried out at Sri Lakshmi Narayana Institute of Medical Sciences, Medical College and Hospital, in collaboration with the Department of Surgery. This research included a total of 93 individuals of various ages who had symptoms of scrotal illness. Patients who did not return for their follow-up appointments were excluded from the research.

CLINICAL ASSESSMENT

A thorough history of clinical symptoms and presenting complaints were collected. To establish the state of scrotal contents, bimanual palpation was used in all patients. As needed, fluctuation, reducibility, and transillumination tests were performed. In cases with suspected testicular cancers, probable metastasis sites were investigated.

RADIOLOGICAL EXAMINATION

Grayscale sonography, CDFI, and PD sonography of scrotal lesions were performed with high-frequency 7 to 12 MHz linear / sector array transducers. When substantial scrotal swellings were present, 5 or 3.5 MHz transducers were required for sufficient penetration. These tests were carried out using a TOSHIBA NEMIO X 2.0. On a monitor extension, picture visualization was acquired. The multiformat camera was used to retrieve the selected photographs in several planes that were saved on the hard drive. The colorful visuals were taken directly from the display and shot.

PREPARATION

There was no need to prepare the patient in any way. Prior to the inspection, pubic hairs were preferred to be shaved. If abdominal sonography was thought to be essential, patients fasted overnight.

INCLUSION CRITERIA:

- All cases with clinical manifestations of testicular pathology.
- Cases of all age groups.

EXCLUSION CRITERIA:

- All cases with lacerated trauma.

TECHNIQUE

The examination took place in a location that provided the patient with enough comfort and privacy. The patient was urged to lie down in a supine position with his legs slightly apart. For support, a towel sling was inserted beneath the scrotum. The penis was lifted and draped on an anterior abdominal wall. The scrotal contents were palpated carefully to begin the examination. A suitable amount of Aquasonic gel was applied to the scrotal area to avoid pressure and discomfort.

The scrotum was scanned from the anterior, lateral, and inferior surfaces in the longitudinal and transverse planes. When necessary, the examination was extended to include the spermatic cord and groyne areas. In cases of varicocele, the examination was carried out in the supine position, using the Valsalva manoeuvre, and then in the upright position to confirm the diagnosis. Masses were photographed and captured in several planes. To avoid pathological masking and optimize contrast resolution in grey scale imaging, special attention was paid to set gains and time gain compensation (TGC). To illustrate small variations in echo texture, a comparison with the contralateral testis proved beneficial.

LABORATORY INVESTIGATIONS

Wherever required, relevant investigations were performed, and a USG-guided FNAC or biopsy was performed. Finally, an investigation was carried out with the goal of correlating clinical results with sonography. CDFI and PD were tested for efficacy in a variety of lesions. The importance of a link between perceptible results and the sonographic examination was emphasized. The sonographic results were associated with histology in 68 individuals. Other patients were sonographically and clinically followed up on.

RESULTS

A total of 93 individuals were assessed utilizing high-resolution grayscale sonography, Color Doppler flow imaging (CDFI), and power Doppler flow imaging (PDFI) for pathological scrotum disorders (PD). These patients varied in age from 3 months to 72 years (Table 1).

TABLE 1: Age distribution of the cases

S.No	Age Groups (Years)	No. of Cases	Percentage (%)
1	0-10	11	11.8
2	11-20	14	15
3	21-30	35	37.6
4	31-40	20	21.5
5	41-50	08	8.6
6	51-60	04	4.3
7	60 and above	01	1
Total		93	100

The significant symptomatology of the patients is shown in Table 2:

TABLE 2: Clinical Symptomatology

S. No	Symptoms	No. of Cases	Percentage (%)
1	Scrotal Swelling	64	44.4
2	Scrotal Pain	46	22.9
3	Empty Scrotal Sac (One Or Both Sides)	4	1.9
4	Trauma	9	4.5
5	Infertility	21	10.5

Scrotal edoema was the most common symptom (36.3 percent), followed by scrotal discomfort (22.9 percent). The majority of the patients experienced several symptoms. Table 3 shows the incidence of intratesticular and extratesticular lesions.

TABLE 3: Incidence of Intratesticular / Extratesticular Lesions

S.no.	Type of Lesion	No. of cases
1.	Intratesticular	15
2.	Extratesticular	77
3.	Intra+Extratesticular	52
Total		144

TABLE 4: Sonographic Features of Malpositioned Testis (N=5)*

Features	No. of cases	Percentage (%)
Position of testis		
• Inguinal	3	60
• Deep Inguinal Ring	2	40
• Other	0	0
Size (compared to normal size)		
• Increased	1	20
• Normal	1	20
• Decreased	3	60
Echo pattern		
• Normal	1	20
• Hypoechoic	3	60
• Hyperechoic	0	0
• Heteroechoic	1	20
Asso. Features		
• Torsion	1	20
• Inguinal Hernia	1	20
• Hydrocele	2	40

*one patient had bilateral undescended testis.

The inguinal canal was the most prevalent place for undescended testis (60 percent). Two patients had associated hydrocele, and one had an inguinal hernia. One patient had torsion of the testes as a complication. The most frequent fluid collection observed was hydrocele. Inflammatory disorders of the scrotum were linked to 31.6 percent of the cases. All cases had unilateral varicoceles. Varicoceles were

present on the left side in 13 cases (86.7%) and on the right in 2 cases (13.3%). All patients were proven surgically to have a varicocele.

Three cases of spermatocele (50%) and six cases of epididymal cysts (75%) affected the head of the epididymis. The contents of spermatocele were echogenic in 83 percent (5/6) of cases, but epididymal cysts were anechoic in all cases. In two cases of spermatoceles and one case of the epididymal cyst, vessels were detected within the septae (Table 5).

TABLE 5: Comparison of Benign and Malignant Testicular Cysts

FEATURES	Benign cysts(n=3)	Malig.cysts(n=4)
Number		
• Single	3	3
• Multiple	0	0
Location		
• Upper pole	2	3
• Lower pole	1	1
• Anterior	2	1
• Posterior	1	3
Cyst walls		
• Smooth, thin	3	0
• Shaggy, thick & poorly marginated	0	4
Unilocular	3	1
Multilocular	0	3
Surrounding soft tissue		
• Normal	3	0
• Abnormal	0	4
Cyst contents		
• Anechoic	3	3
• Echogenic	0	1
CDFI AND PD FEATURES		
• Vascularity of the testis	Normal	Hypervascular
• Vascularity of the cysts	avascular	Avascular

In this investigation, benign cysts were solitary, unilocular, and had smooth thin walls. Malignant cysts, on the other hand, had shaggy, thick, and weakly marginated walls in all cases and were multilocular in 75% of them. In instances of benign cysts, normal testicular parenchyma was evident surrounding them, whereas tumour parenchyma was apparent around malignant cysts.

DISCUSSION

The goal of this study was to compare the multidimensional data generated by high-resolution grayscale sonography and Color Doppler flow imaging in the diagnosis of scrotal disease. This study comprised 93 individuals who had indications of a lesion in the scrotal tissues. Young guys were more likely to have scrotal diseases. Scrotal edoema was the most common complaint, followed by scrotal discomfort [16–18]. A total of 144 pathological lesions were found in the 93 individuals who took part in the research. In 5.3 percent of our patients, cryptorchidism was discovered. The right side of the body was heavily affected. The inguinal canal was the most prevalent site for cryptorchid testes in our patients (60 percent). Sixty percent of the testes were homogeneous and hypoechoic. One example involved the torsion of undescended testes. On the contralateral side, two testes had grade I vascular signals, whereas the other had grade II.

CONCLUSION

In the context of clinical and laboratory data, different parameters in scrotal illnesses have been researched and analyzed sonographically (grayscale, CDFI, and PD). The following conclusions may be drawn from this research: Scrotal illnesses were seen in all age categories, with young guys having the highest prevalence. The most prevalent clinical signs were scrotal swellings followed by scrotal discomfort. The anatomical characteristics and vascularity of scrotal lesions could be consistently defined using high-resolution sonography with colour Doppler flow imaging (CDFI) and power Doppler (PD). Sonography was quite accurate in determining whether a scrotal tumour was solid or cystic. In this investigation, 100 percent accuracy was attained in this regard. Sonography was helpful in determining if a scrotal abnormality was

intra or extra-testicular. This was crucial because practically all additional testicular diseases are harmless. In 98 percent of our situations, this was doable. In the diagnosis of hydrocele, sonography was found to be 100 percent sensitive. It also aided in determining the integrity and size of testes in situations with big hydrocele, which is a tough clinical task. Idiopathic hydrocele might be distinguished from those caused by trauma or infections using sonography. High-resolution sonography might be employed as a first-line inquiry in the evaluation of scrotal diseases, according to the findings of this study.

FUNDING: No funding sources

ETHICAL APPROVAL: The study was approved by the Institutional Ethics Committee.

CONFLICT OF INTEREST: The authors declare no conflict of interest.

ACKNOWLEDGMENTS

The encouragement and support from Bharath University, Chennai is gratefully acknowledged. For provided the laboratory facilities to carry out the research work.

REFERENCES

1. Evans JP, Snyder HM. (1977). Idiopathic scrotal edema. *Urology*; 9: 549–551.
2. de Jesus LE. (2008). Acute Scrotum BT - Normal and Abnormal Scrotum. In: Fahmy MAB (ed). Cham: Springer International Publishing, pp. 299–318.
3. Piçarro C, Lima DX. (2007). Common Acquired Diseases of the Scrotum BT - Normal and Abnormal Scrotum. In: Fahmy MAB (ed). Cham: Springer International Publishing, pp. 319–339.
4. Knight PJ, Vassy LE.(1984). The diagnosis and treatment of the acute scrotum in children and adolescents. *Ann Surg* ; 200: 664–673.
5. Valentino M, Bertolotto M, Derchi L, et al. (2011). Role of contrast enhanced ultrasound in acute scrotal diseases. *Eur Radiol*; 21: 1831.
6. Miskin M, Bain J.(1974). B-mode ultrasonic examination of the testes. *J Clin Ultrasound*; 2: 307–311.
7. Ragheb D, Higgins Jr JL. (2002). Ultrasonography of the Scrotum. *J Ultrasound Med*; 21: 171–185.
8. Schwarze V, Marschner C, Rübenthaler J, et al. (2020). Overview of ultrasound applications for assessing scrotal disorders. *J Ultrasound Med*; 39: 1047–1056.
9. Patelli E, Gelosa M, Mantovani F, et al. (1998). Surgery in day hospital: scrotal surgery]. *Arch Ital di Urol Androl organo Uff [di] Soc Ital di Ecogr Urol e Nefrol*; 70: 121–125.
10. Frush DP, Applegate K. (2004). Computed tomography and radiation: understanding the issues. *J Am Coll Radiol* ; 1: 113–119.
11. Rehani MM, Berry M. (2000). Radiation doses in computed tomography. *BMJ* ; 320: 593 LP – 594.
12. Kühn AL, Scortegagna E, Nowitzki KM, et al. (2016). Ultrasonography of the scrotum in adults. *Ultrason (Seoul, Korea)* ; 35: 180–197.
13. Hamm B. (1994). Sonography of the testis and epididymis. *Andrologia*; 26: 193–210.
14. Anand SS, Mukherjee PB, Kumar PG, et al. (2008). Scrotal Scintigraphy in a Case of Acute Scrotum. *Med journal, Armed Forces India*; 64: 92–94.
15. Erden MI, Özbek SS, Aytaç SK, et al. (1993). Color-Doppler Imaging in Acute Scrotal Disorders. *Urol Int*; 50: 39–42.
16. Zhang H, Li J, Saravanan KM, et al. (2021). An Integrated Deep Learning and Molecular Dynamics Simulation-Based Screening Pipeline Identifies Inhibitors of a New Cancer Drug Target TIPE2. *Front Pharmacol*; 12. Epub ahead of print . DOI: 10.3389/fphar.2021.772296.
17. Saravanan KM, Selvaraj S. (2017). Dihedral angle preferences of amino acid residues forming various non-local interactions in proteins. *J Biol Phys* ; 43: 265–278.
18. Zhang H, Shao X, Peng Y, et al. (2019). A novel machine learning based approach for iPS progenitor cell identification. *PLoS Comput Biol*; 15. Epub ahead of print. DOI: 10.1371/journal.pcbi.1007351.

CITATION OF THIS ARTICLE

A K Gupta, S S Shetty, G. Jayalakshmi. Scrotal Pathology Evaluation Using High-Resolution Sonography Imaging. Bull. Env.Pharmacol. Life Sci., Vol 12 [1] December 2022 : 15-19