



## ORIGINAL ARTICLE

# The Laboratory Scale Evaluation of Multiple pH ranges on *Spirulina platensis* culture in the Production of Dry Biomass, Chlorophyll, Phycocyanin, & Carotenoids

Ehsan Hosseini<sup>1,4</sup>, Mahboobe Mohamadami<sup>2</sup>, Mohsen Akbari<sup>3</sup>, Faranak Nejati<sup>4</sup>, Manoucher Mazdapour<sup>5</sup>, Mozhgan Ghasemian<sup>6</sup>, Armin Bahmani\*<sup>2</sup>

<sup>1</sup>Department of Bacterial Vaccine and Antigen Production, Pasteur Institute of Iran, Tehran, Iran

<sup>2</sup>Department of Biology, Islamic Azad university of Karaj, Iran

<sup>3</sup>Department of Biology, Islamic Azad university of Ghom, Iran

<sup>4</sup>Department of Biology, Islamic Azad university of Damghan, Semnan, Iran

<sup>5</sup>Department of Biology, Pharmaceutical Sciences Branch, Islamic Azad University, Tehran, Iran

<sup>6</sup>Department of Biology, Islamic Azad university of Falavarjan, Isfahan, Iran

Email: Armin.ba84@yahoo.com

### ABSTRACT

*Spirulina* is a complete food resource of Chlorophyll, Phycocyanin, & Carotenoids. It is also has an application as a natural dye in food industry, cosmetic and pharmaceutical industry. In this study, the effects of a wide range of pH were evaluated on the growth of *Spirulina platensis* on various media such as, ZM, SWMX2, NRCM, CFTRI, & Wastewater. In addition, the pH modification was also done to evaluate the rates of dry weight, phycocyanin, & chlorophyll production. The *Spirulina platensis* cultivation was done on ZM, SWMX2, NRCM, CFTRI, & Wastewater media and the extraction of chlorophyll, phycocyanin, and carotenoid were done by ultrasonication, homogenation, & acetone methods, then all products were analyzed by HPLC. In this study, the pH ranges of 8-10.8 were used in the *Spirulina platensis* cultivation. The dry weight, chlorophyll, and phycocyanin concentration was in ZM medium with pH=9.6, 9, and 9.6 was 2.28g/100ml, 12.83 mg/g, & 44.27 mg/g, respectively. The highest concentration of carotenoid was in CFTRI medium with pH= 10.2, as 3.13 mg/g. All the media used in this study had an acceptable growth and production. Although, the pH range of 9-10.2 had the highest efficacy of production and growth on *Spirulina platensis*.

**Keywords:** *Spirulina platensis*, Dry Weight, Chlorophyll, Phycocyanin, Carotenoid.

Received 01.10.2014

Revised 01.11.2014

Accepted 08.12.2014

### INTRODUCTION

Since centuries cyanobacteria have been receiving increasing interest due to their potential to produce a diverse range of chemicals and biologically active compounds, such as vitamins, carotenoid pigments, proteins, lipids and polysaccharides (1). For exploration of these potentials of cyanobacteria it should be cultivated in commercial way. Globally researchers are trying to produce microalgae/cyanobacteria commercially [2, 3]. Yet very little or primary information is available on detailed design criteria, location selection, scaling considerations, or constrains involved in large scale cultivation. *Spirulina* is a planktonic photosynthetic filamentous cyanobacterium that forms massive populations in tropical and subtropical bodies of water which have high levels of carbonate and bicarbonate and alkaline pH values of up to 11. *Spirulina* from Chad Lake in Africa and Texcoco Lake in Mexico have been harvested as a source of food [4]. *Spirulina* has been studied for single cell protein (SPC) [5], vitamins, minerals, proteins and polyunsaturated fatty acids (gamma-linolenic acid) [6], therapeutic properties [2, 3], antioxidant activity (7). Several cultivation methods like; open ponds [8], tubular photobioreactors (9), inclined glass panels (10) have been tried. Cost and composition of cultivation media along rate of the algae as challenging factors for commercially viable production. Different media have been tried for cultivation of *Spirulina* such as Zarrouk's media [11], SWMX2 media [12], NRCM media [13] CFTIR media, Wastewater media [14]. Recently, it has been focused on valuable products such as chlorophyll, phycocyanin, and

carotenoids. This study has been done to evaluate the effects of various pH ranges of different selected media on the production rate of dry biomass weight, chlorophyll, phycocyanin, and carotenoids.

## MATERIAL AND METHODS

### Isolation & Mass Cultivation of Spirulina

The water sample was inoculated into the Zarrouk medium for isolation of *Spirulina* and incubated in a growth chamber with light flux of 12:12 h, light intensity of 4Klux, in 30°C and pH=9. The purity of the culture was ensured by repeated inoculation, and identification was accomplished by determining cellular morphology observed by using light microscope.

### Culture medium:

Five media were included in this study. For this experiment Zarrouk's media (ZM), SWMX2 medium, NRCM, CFTRI medium, Wastewater Medium, were chosen (the composition of all media are described in Table 1, respectively). Prepared 1000ml each medium in 1000ml flask and inoculate 50ml of inoculums in each medium. After 28 days the biomass of *Spirulina platensis* was harvested and the dry weight was determined.

### Extraction of chlorophyll:

1g of wet spirulina biomass was taken in a beaker and about 10ml of aqueous acetone solution (Sigma,USA) was added, closed tightly and placed in a dark box overnight. Then sonicated by ultrasonicator for 20 second at the 5 setting. The extract was then brought to a volume of 13 ml with aqueous acetone solution, which was closed tightly and kept again in the dark box overnight in a cold room. The next day the extract was transferred to a centrifuge tube and mixed well. Then the extract was clarified by centrifugation for 20 minutes at 1000 rpm. The pellet was discarded and supernatant was separated which contained the chlorophyll pigments [15].

### Phycocyanin extraction:

The C-phycocyanin was extracted from fresh biomass by the following procedure. Fresh biomass was homogenized with 50mM sodium phosphate buffer, the homogenate was subjected to alternate freezing and thawing( 3 to 4 cycles) and centrifuged at 5000rpm for 10 minutes. The phycocyanin content was estimated.

### Carotenoid extraction:

Total carotenoids were extracted by cell lysis (sonication) and homogenization in 90 percent acetone as described by [11]. The final extract was measured at A450 nm against acetone as blank. As for  $\beta$ -carotene, the cells were homogenised using acetone and petroleum ether and the extracts were pooled and measured at A453 nm against petroleum ether as blank [16].

### HPLC analysis:

The chemical constituents of the *S.platensis* extracts were identified by HPLC method reported by [17]. Dionex Summit IV HPLC system consisted of a Dionex P680 dual gradient pump, an ASI-100 auto-sampler equipped with a 20- mL loop and PDA- 100 photodiode array detector were used. A reversed phase column C18 (250 x 4.6 mm, 5 mm partials) was used. The mobile phase was a mixture of solvent A (methanol/ammonium acetate 0.1 N; 7:3, v/v) and solvent B (pure methanol) at rate of 0.9 ml min<sup>-1</sup> as a step gradient, lasting 35 min, which started from 25% B, changing at 50% in 1 min, rising up to 100% B at 10 min. Then, the mobile phase composition was kept constant until the end of the analysis. Total acquisition time was 35 min. The temperature was set at 25°C. The identification of the peaks was performed, when possible, using standards. When no standards were available, tentative identification was done based on UV-Vis spectra characteristics and compared with that data appearing in the literature.

### Statistical analyses:

The data recorded in triplicate for the parameters in various strains were subjected to ANOVA (analysis of variance).

## RESULTS

### The Results of Light Microscopy:

The microscopy result of *Spirulina platensis* cultivation in the media used in this study, which was seen as a spiral-shaped cyanobacterium in the microscopy field.




Fig 1: Spirulina platensis

The effects of pH on obtaining the highest dry weight could be seen on tables 2,3& 4. The cultivation done on ZM, SWMX2, NRCM, CFTRI, and Wastewater at the pH ranges 9.6, 8.6, 8.8, 9, 9.8 had the highest concentration of 2.28±0.13, 1.16±0.52, 0.95±0.94, 1.75±0.04 g/l, respectively.

Table.1 indicates the contents of all media used in this study. The data are mentioned as g/l.

	ZM	SWMX2	NRCM	CFTRI	Wastewater
NA HCO <sub>3</sub>	16/8	0/2	8	40	8
K <sub>2</sub> HPO <sub>4</sub>	0/5	0/016	0/5	0/5	1
NaNO <sub>3</sub>	2/5	2	2/5	1/5	1/5
K <sub>2</sub> SO <sub>4</sub>	1	0/5	1		
NaCL	1	5	1		
MgSO <sub>4</sub> .7H <sub>2</sub> O	0/2	0/16	0/2		
CaCL.2H <sub>2</sub> O	0/04	0/04			
FeSO <sub>4</sub> .7H <sub>2</sub> O	0/01	0/01	0/01		
EDTA	0/08	0/01			

Tables.2 to.4 indicate The effects of pH on obtaining the highest dry weight could be seen on tables 2,3 & 4. The cultivation done on ZM, SWMX2, NRCM, CFTRI, and Wastewater at the pH ranges 9.6, 8.6, 8.8, 9, 9.8 had the highest concentration of 2.28±0.13, 1.16±0.52, 0.95±0.94, 1.75±0.04 g/l, respectively.

Table 2 :Effect of PH on biomass (dry wt) of S.platensis in ZM,SWMX2,NRCM,CFTRI,Wastewater media.

Medium	pH				
	8	8.2	8.4	8.6	8.8
ZM	0.86±0.36	0.91±0.62	0.99±0.07	1.2±0.79	1.24±0.02
SWMX2	0.97±0.32	0.99±0.66	0.1±0.96	1.16±0.52	1.05±0.25
NRCM	0.53±0.18	0.61±0.09	0.66±0.78	0.88±0.49	0.95±0.94
CFTRI	0.72±0.38	0.77±0.57	0.87±0.67	1.01±0.97	1.16±0.56
Wastewater	0.61±0.58	0.55±0.62	0.79±0.53	0.82±0.64	0.92±0.67

Mean ± SD in each column with different superscripts indicating significance (P < 0.05)

Table 3: Effect of PH on biomass (dry wt) of S.platensis in ZM, SWMX2, NRCM,CFTRI,Wastewater media.

Medium	PH				
	9	9.2	9.4	9.6	9.8
ZM	1.84 ±0.46	1.92±0.53	2.13±0.49	2.28±0.13	1.97±0.063
SWMX2	0.84±0.15	0.87±0.48	0.91±0.25	0.95±0.61	0.72±0.27
NRCM	1.42±0.057	1.53±0.98	1.65±0.19	1.91±0.041	1.54±0.18
CFTRI	1.75±0.04	1.46±0.57	1.44±0.12	1.26±0.58	1.17±0.01
Wastewater	0.26±0.16	0.28±0.69	0.49±0.76	0.68±0.38	0.94±0.47


Mean ± SD in each column with different superscripts indicating significance (P < 0.05)

Table 4: Effect of PH on biomass (dry wt) of *S.platensis* in ZM,SWMX2,NRCM,CFTRI,Wastewater medium.


Medium	pH				
	10	10.2	10.4	10.6	10.8
ZM	1.26±0.12	1.42±0.07	1.11±0.95	0.96±0.66	0.89±0.57
SWMX2	0.73±0.36	0.66±0.38	0.77±0.97	0.64±0.28	0.41±0.32
NRCM	0.55±0.69	0.49±0.67	0.48±0.55	0.33±0.01	0.28±0.96
CFTRI	0.81±0.36	0.79±0.36	0.92±0.65	0.96±0.91	0.84±0.39
Wastewater	0.56±0.25	0.72±0.36	0.76±0.49	0.37±0.95	0.16±0.09

Mean ± SD in each column with different superscripts indicating significance ( $P \leq 0.05$ )


Graph.1, Graph.2, Graph.3, Graph .4, Graph.5 all indicate the rate of chlorophyll, phycocyanin, carotenoid production in the media ZM, SWMX2, NRCM, CFTRI, & Wastewater, in the pH range of 8-10.8, respectively.




Graph.2




Graph.3



Graph.4



Graph.5



As it could be seen, in order to obtain the concentrations of chlorophyll, phycocyanin, and carotenoid, the pH is essential in various media, which as shown in graph 1-5, had the highest value in ZM medium. The results for the highest chlorophyll, phycocyanin and carotenoid in this medium at the pH= 9.6-9.6-8.4 are 12.83 mg/g, 44.27 mg/g, and 3mg/g, respectively. The highest concentration of chlorophyll, phycocyanin, and carotenoid was in SWMX2 medium with the pH range of 10.4-8.6-9.8, and the results were 2.62 mg/g, 15.23 mg/g, and 3.4 mg/g, respectively. The highest concentration of chlorophyll, phycocyanin, and carotenoid was in NRCM medium with the pH =9.2-9.6-10.4, and the results were 12.1 mg/g, 35.99 mg/g, 3.58 mg/g, respectively. The highest concentration of chlorophyll, phycocyanin, and carotenoid was in CFTRI medium with the pH range of 8.8-9-10.2, and the results were 12.31 mg/g, 7.98 mg/g, 3.13 mg/g, respectively. The highest concentration of chlorophyll, phycocyanin, and carotenoid was in Wastewater with the pH range of, 9.2-9.8-10.2 and the results were 2.92 mg/g, 12.27 mg/g, 2.46 mg/g, respectively.

**DISCUSSION**

*Spirulina platensis* was grown at various pH ranges (8-10.8) in flask culture and monitored and expressed in term of dry weight (Table 1-3). The maximum bulk density about 2.28 g/1000ml was noticed when the pH of culture medium was, maintained at 9.6 with medium volume 1000 ml flask. The maximum bulk density was attained on 28<sup>th</sup> day after the inoculation of culture in medium. The increase in the production of *Spirulina platensis* could have been due to the availability of mire space, oxygen and light to the culture flask. Earlier results also demonstrated that optimum pH for maximum growth of *Spirulina platensis* was 9 to 9.5 ranges [18]. *Spirulina platensis* is considered to be an alkalophilic organism by nature (19). The Chlorophyll a content is also maximum in pH 9, 12.83 mg/g of dry weight. Similar studies have also been done by other scientists on cyanobacteria [20, 21]. Phycocyanin content is also maximum in pH 9.6, 44.27 mg/g of dry weight. Similar studies have also been done by various workers of cyanobacteria [22]. The Carotenoid content is also maximum in pH 10.2, 3.13 mg/g of dry weight. Similar studies have also been done by various workers of cyanobacteria [23].

**ACKNOWLEDGEMENTS**

The authors of this study would like to acknowledge all institutes for supporting and to providing the funding of this project, especially the deputy of the research & production complex, Dr. Hooman Kaghazian and the department head of bacterial vaccines & antigen production of pasteur institute of Iran, Mr. Mehdi Nejati. We would also like to thank all consultants for their precious assistance to this project, especially Dr. Soltani, Dr. H. Ahmadi, Dr. Dadkhalipour, & Mr. M.H. Hedayati.

**REFERENCES**

- Zhang, X.-W., Zhang, Y. M., Chen, F. (1999). Application of mathematical models to the determination optimal glucose concentration and light intensity for mixotrophic culture of *Spirulina platensis*. *Process Biochem.* 34: 477-481.
- Belay, A., Ota, Y., Miyakawa, K., Shimamatsu, H., (1993). Current knowledge on potential health benefits of *Spirulina*. *Journal of Applied Phycology.* 5: 235-241.
- Belay A., (1997). Mass culture of *Spirulina* outdoors – the Earthrise experience. In: Vonshak A (ed) *Spirulina platensis* (Arthrospira): physiology, cell-biology and biotechnology. Taylor & Francis, London, pp 131-158.
- Vonshak, A., (1997). *Spirulina platensis* (Arthrospira). Physiology, Cell biology and Biotechnology. Taylor & Francis, London.
- Anupama, P.R., (2000). Value-added food: single cell protein. *Biotechnology Advances.* 18: 459-479.
- Miranda, M.S., Cintra, R.G., Barros, S.B.M., Filho, J.M. (1998). Antioxidant activity of the microalga *Spirulina maxima*. *Brazilian J. of Medical and Bio. Res.* 31: 1075-1079.
- Estrada, J.E., Bescós, P., Villar Del Fresno, A.M. (2001). Antioxidant activity of different fractions of *Spirulina platensis* protein extract. *Farmaco* 56: 497-500.
- Lee YK. 1997. Commercial production of microalgae in the Asia-Pacific rim. *J. Appl. Phycol.* 9: 403-411.
- Torzillo G., Pushparaj B., Bocci F. (1986). Production of *Spirulina* biomass in closed photobioreactors. *Biomass.* 11: 61-74.
- Hu Q, Guterman H, Richmond A. 1996. A flat inclined modular photobioreactor (FIMP) for outdoor mass cultivation of photoautotrophs. *Biotechnol. Bioeng.* 51: 51-60.
- Zarrouk, C. 1966. Contribution to the study of cyanophyceae. Influence of various physical factors et chimiques sur la croissance et la photosynthèse de *Spirulina maxima*. PhD Thesis, University of Paris, Paris.
- Facundo J. Márquez-Rocha, Phycocyanin Production In Seawater Culture of *Arthrospira maxima*, *Ciencias Marinas*, Vol. 26, No. 4.
- Rafiqul, I.M., A. Hassan, G. Sulebele, C.A. Orosco, P. Roustaiian and Jalal, C.A. (2003). Salt stress culture of blue green algae *Spirulina fusiformis*. *Pak. J Biol. Sci.* 6 (7): 648-650.
- Anaga, A.; Abu, G. O. A laboratory-scale cultivation of *Chlorella* and *Spirulina* using waste effluent from a fertilizer company in Nigeria. *Bioresource Technology*, v. 58, p. 93-95, 1996.
- Nelson, D.H. 1960. Improved chlorophyll extraction method. *Science*, 132:351-352.
- Govindaraju, P., T. Devaki and Subash Chandra Bose, M. 2001. Estimation of lycopenes and carotenes. In: *Practical plant Biochemistry*. Rasi publications, 74-75.
- Mendiola JA, Marin FR, Hernandez SF, et al., (2005). Characterization via liquid chromatography coupled to diode array detector and tandem mass spectrometry of supercritical fluid antioxidant extracts of *Spirulina platensis* microalga. *J. Separa. Sci.* 28: 1031-1038.
- Belkin, S. and Boussiba, S. (1971). Resistance of *Spirulina platensis* (Cyanophyta) to high pH values. *Plant cell Physiol.* 32:953-9589.
- Grant, W.D., Mwatha, W.E. and Jones, B.E. (1990). Alkalophiles: ecology, diversity and application. *FEMS Microbiol rev.* 75:225-270.
- Carvalho, J.C.M., Sato, S., Moraes, I. DE O. and Pelizer, L.H. (2002). *Spirulina platensis* growth estimation by pH determination at different cultivation conditions. *Electronic Journal of Biotechnology.* 5(3):251-257.
- Kim C.J., Jung, Y.H. and OH, H.M. (2007). Factors indicating culture status during cultivation of *Spirulina* (Arthrospira) *platensis*. *The Journal of Microbiology.* 45 (2):122-127.
- Xianhai Zeng, Michael K. Danquah, Shiduo Zhang, et al, (2012). Autotrophic cultivation of *Spirulina platensis* for CO<sub>2</sub> fixation and phycocyanin Production, *Chemical Engineering Journal* 183; 192- 197.
- K.Sujatha and P.Nagarajan, (2013). Optimization of growth conditions for carotenoid production from *Spirulina platensis* (Geitler), *Int. J. Curr. Microbiol. App. Sci.* 2(10): 325-328.

**CITATION OF THIS ARTICLE**

Ehsan H, Mahboobe M, Mohsen A, Faranak N, Manoucher M, Mozghan G, Armin B. The Laboratory Scale Evaluation of Multiple pH ranges on *Spirulina platensis* culture in the Production of Dry Biomass, Chlorophyll, Phycocyanin, & Carotenoids. *Bull. Env. Pharmacol. Life Sci.*, Vol 4 [2] January 2015:13-18