



Cytotoxic, antimicrobial and antioxidant activity of *Viburnum grandiflorum* leaf extracts: growing wild in PirPanjal Range of Kashmir Valley

AbRouf Wani^{1*}, Sugandha Singh²

¹Department of Botany, Madhyanchal Professional University, Ratibad, Bhopal, M.P., India.

²Dean Faculty of Sciences and IT, Madhyanchal Professional University, Ratibad, Bhopal, M.P., India.

*Corresponding author: AbRoufWani. Email ID: roufwani232@gmail.com

ABSTRACT

The main aim of this research work was to evaluate the cytotoxic effects, antibacterial and antioxidant activity of different solvent extracts of *Viburnum grandiflorum* leaves. The extraction of the leaves was done with ethanol, ethyl acetate, n-hexane and water using Soxhlet apparatus. The agar well diffusion method was used to determine the antibacterial activity of the extracts. The antioxidant activity was measured by using DPPH scavenging assay and hydrogen peroxide reduction method. The cytotoxic studies were carried out by MTT assay using A549 lung cancer cells. Results showed remarkable antibacterial activity of the four extracts against the selected bacteria but higher inhibition and lower MIC values were shown by ethanolic extract with 21.7 ± 0.37 mm and 0.1 mg/ml; 29.5 ± 0.45 mm and 0.037 mg/ml; 30.5 ± 0.21 mm and 0.07 mg/ml, respectively. Similarly, all the extracts showed significant antioxidant and cytotoxic activity in a concentration-dependent manner but the ethanolic extract was the leading one with higher antioxidant activity and a proliferation inhibition of upto 95% in A549 lung cancer cells. Therefore, our findings showed that the leaves of *Viburnum grandiflorum* possess significant antibacterial, antioxidant and cytotoxic ability. This indicates that more emphasis should be given on finding the bioactive constituents of the plant responsible for its medicinal activities.

Keywords: *Viburnum grandiflorum*, extract, antioxidant, antibacterial, cytotoxic

Received 11.03.2022

Revised 26.06.2022

Accepted 07.07.2022

INTRODUCTION

The Genus *Viburnum* under the family of Adoxaceae comprises of over 200 small trees and shrubs [1]. These plants are predominantly found in the Northern Hemisphere with temperate climatic zone and the tropical mountains of Tasmania and Australia. Significant diversities of these plants are seen in China and Himalayan mountains [2]. In the earlier times the health benefits and therapeutic value of *Viburnums* were well understood. The plant *Viburnum grandiflorum* grows extensively in wild in PirPanjal mountain range of Kashmir Valley. The *Viburnum grandiflorum* Wall ex D.C. has a very rich therapeutic value and is used for the treatment of malaria, wounds and as a diuretic [3]. *Viburnum grandiflorum* is used by indigenous people as a purgative, abdominal pain, diuretic and antimalarial [4,5]. It is used to treat typhoid, toothache, respiratory disorders, whooping cough, upset stomach, wounds and as an anesthetic [6-8]. The noteworthy contribution of Uddin *et al.* in the field of conventional medicinal system concerning the medicinal importance of *Viburnum grandiflorum* as an antipyretic agent against the malaria and typhoid [9]. Their contribution has provided scientific justification for different pharmaceutical role of *Viburnum grandiflorum* like antipyretic, anti-nociceptive, and anti-inflammatory, with respect to its phytochemical composition. Moreover, the *Viburnum* species is used to treat several health conditions like antibacterial, antioxidant, anti-diabetic, rheumatoid and diarrhea [10-12].

The lung cancer is a lethal respiratory condition which remains the most common form of cancer-related deaths, globally. The lung cancer is hence ranks first in terms of death rate worldwide. The overall lung cancer related deaths are very high than that of collectively caused by brain, colorectal, breast, prostate cancers [13]. It is estimated that the year 2021 will encounter over 235,760 new cases and about 131,880 deaths alone in United States [14]. The non-small cell lung cancer is the most frequently prevailing form of lung cancer, accounting for more than two-thirds of the cases and majority (84%) of the patients are diagnosed at advanced stage of the disease [15]. Therefore, lung cancer is a foremost lethal form of cancer globally, that needs to be addressed with novel therapies and chemopreventive agents. Natural products

are one of the leading strategies to tackle this condition because of higher diversities in chemical species and healthy biological profiles [16]. Antioxidants are the health benefiting species, which protects body against oxidative stress caused by the generation of free radical species [17]. The medicinal plants bear phytochemicals like polyphenols show very high antioxidant capacity. Furthermore, the medicinal plants have served humanity with a number of antibiotic drugs like penicillin.

Therefore, the present study was designed by keeping in view the medicinal benefits of *Viburnum grandiflorum*. We investigated the antioxidant, cytotoxic and antimicrobial activity of *Viburnum grandiflorum* leaf extracts growing in PirPanjhal mountain range of Kashmir Valley.

MATERIAL AND METHODS

Collection of the plant material

The plant material was collected in August, 2020, from Yusmarg area of PirPanjhal mountain range in Kashmir Valley. The plant was authenticated by Dr. Kanchan Yadav, Professor Department of Botany, Madhyaanchal Professional University, Ratibad, Bhopal, India.

Extract preparation

The leaves of the plant *Viburnum grandiflorum* were collected and washed under running water. Washed leaves were dehydrated under shady condition with a temperature of 31 ± 2 °C for one week. The dried leaves were crushed to fine powder using motor and pestle and then stored in airtight plastic bags. The 100g of the powder was extracted with different solvents including 70% ethanol, ethyl acetate, *n*-hexane and water, using Soxhlet extractor for 12h. The extract then was placed in a rotatory evaporator for elimination of the moisture and remaining solvent. Afterwards, different solvent concentrations viz 30, 60, 120, 240, and 480 µg/ml were prepared using DMSO as vehicle control.

Microbial culture

The antibacterial activity of the leaf extracts of *Viburnum grandiflorum* was measured against three bacterial strains including *M. luteus*, *S. aureus*, and *E. coli*. The microbial cultures were procured from the Microbial Type Culture Collection, Institute of Microbial Technology, Chandigarh, India. The bacterial viability was maintained by culturing each strain in Muller-Hinton agar slants every two weeks. The experimental settings were completely in agreement with the provided protocol by CLSI Protocol requirements.

Antibacterial analysis of *Viburnum grandiflorum* leaf extracts

The agar well diffusion method was applied to determine the antibacterial activity of the leaf extracts of *Viburnum grandiflorum* including ethanolic, ethyl acetate, *n*-hexane, and aqueous extracts. The bacterial strains were cultured for 12h in absence of light at 37°C within Mueller-Hinton Broth. About 100 µl from each bacterial culture was taken and inoculated within the molten Muller-Hinton Agar. This was followed by homogenizing and transference to 90mm petri dishes (sterilized) which was allowed to settle down in laminar air flow. The 5mm diameter standard cork barrier was used to create wells, to which 50 µl of different extracts of *Viburnum grandiflorum* leaves in DMSO were added. The DMSO was taken as negative control and 10 µg/disc of Ampicillin was taken as positive control. Each petri dish was protected by a laboratory film to avoid evaporation followed by the incubation for 24 h at 37°C. A standard scale was finally used to mark the zones of inhibition (mm) to the nearby size. The experimental procedures for each extract were repeated in triplicates and data was expressed as mean \pm SD.

Determination of minimum inhibition concentration

The broth dilution method was used to determine the MIC of the extracts of *Viburnum grandiflorum*. The different concentrations ranging from 0-480 µg/ml of ethanolic, ethyl acetate, *n*-hexane and aqueous extracts were made by dilution with Muller-Hinton broth. The turbidity of the bacterial strains was set at 0.5 McFarland standard turbidity. All the samples were shaken vigorously followed by inoculation with test bacterial suspension of 50 µl maintaining 2×10^4 CFU/ml. This mixture was then incubated for 24h at 37°C and finally the concentration at which no sign of apparent bacterial growth is seen was taken as MIC value.

Antioxidant activity

DPPH radical scavenging assay

The fresh DPPH solution was prepared with methanol by adding 2.8 mL of 100 µM DPPH methanol. Afterwards, different concentrations of the extracts viz 30, 60, 120, 240, and 480 µg/ml were prepared with 0.2mL of methanol and added to this mixture. These mixtures were then subjected to incubation for half an hour at room temperature. The DPPH free radical has a tendency of one free electron and higher absorbance at 517 nm. The DPPH solution shows deep purple coloration, which gets transformed to golden yellow on addition of hydrogen atom from resultant donor. The fading/discoloration is directly a signifying characteristic of its reduction and concentration and antioxidant activity of the extracts tested. Post incubation, the mixture was forwarded to absorbance measurements at 517 nm, with methanol

serving as blank and methanol and DPPH solution deprived of the extract concentrations serves as control. Each of the experiment was performed in triplicates and ascorbic acid was used as reference control. The percentage inhibition of DPPH radicals was calculating according to the equation:

$$\%Inhibition = \frac{Absorbance\ of\ control - Absorbance\ of\ sample}{Absorbance\ of\ control} \times 100$$

Hydrogen peroxide reduction/scavenging assay

In brief, the various experimental concentrations of extracts and reference control viz 30, 60, 120, 240, and 480 µg/ml were prepared in DMSO. Independently, a mixture of 40 mM of H₂O₂ was prepared in phosphate buffer of 0.1 M and pH 7.4. The extract and H₂O₂ mixtures were mixed together and incubated for 20 min. The absorbance was recorded at 230 nm to detect the radical scavenging activity. The percentage inhibition was calculated using following equation:

$$\% Inhibition = (Absorbance\ of\ reaction\ mixture / Absorbance\ of\ sample) \times 100$$

Cell culture and conditions

The lung cancer cell line A549 was obtained from American Type Culture Collection (Manassas, United States). The A549 cells were placed in DMEM maintaining 10% fetal bovine serum and penicillin and streptomycin as antibiotics. This cell culture was placed under humidified environment inside a 5% CO₂ incubator at 37°C.

MTT assay

The evaluation of cytotoxic activity of *Viburnum grandiflorum* leaf extracts against the lung cancer A549 cells was executed by implementing MTT assay. In brief, A549 cell line was pre-cultured and then subjected to treatment with separate concentrations of ethanolic, ethyl acetate, *n*-hexane and aqueous extracts ranging from 0-480 µg/ml in 96-well plates. The extract treated A549 cells were then incubated for 48 h at 37°C and then washed with PBS. Following PBS washing, cells were added with MTT stock solution. The MTT has a tendency to get reduced by viable cells to insoluble formazan crystals. Therefore, the generation of formazan crystals is directly proportional to the viable cells in MTT assay. Then, the formazan crystals formed were dissolved in the DMSO (100 mL) and subjected for colorimetric analysis using a microplate reader.

Statistical analysis

The individual experiments were executed in triplicates and data was shown as mean ± SD. One way ANOVA was used for data analysis followed by statistical comparisons by student's t-test using SPSS software version 15.0. The p-value of less than 0.05 was taken as statistically significant figure.

RESULTS AND DISCUSSION

Antioxidant activity of the leaf extracts of *Viburnum grandiflorum*

Medicinal plants are very rich in bioactive scaffolds which act as antioxidants or free radical scavengers in biological systems [18]. The free radical or oxidative species are very high in reactivity so that they interfere with everything that comes in contact with them. In biological systems, these oxidative species are responsible for damaging normal functions by interfering with key biomolecules and enzymes [19]. Moreover, these highly reactive species cause DNA damage and hence take part in carcinogenesis. Therefore, monitoring and antioxidants are of high demand for health promotion. Herein, we investigated the antioxidant effects of different solvent extracts of *Viburnum grandiflorum* leaves including ethanolic, ethyl acetate, *n*-hexane and aqueous extracts.

DPPH scavenging activity

To test the antioxidant potential of different solvent extracts of *Viburnum grandiflorum* leaves, we performed DPPH scavenging assay. The DPPH scavenging activity of the extracts was taken at different concentrations ranging from 30-480 µg/ml. The results showed promising antioxidant capacity of the extracts in a concentration-dependent manner (Figure 1). The ethanolic extract showed significant DPPH scavenging followed by the ethyl acetate, aqueous and *n*-hexane extracts in comparison to that of reference control ascorbic acid. On higher extract concentrations (480 µg/ml), the ethanolic, ethyl acetate, aqueous and *n*-hexane extracts showed percentage DPPH inhibition of 83%, 79%, 59%, and 53%, respectively. Therefore, based on DPPH scavenging assay it is observed that the leaves of *Viburnum grandiflorum* possess strong DPPH scavenging activity.




Figure 1: The DPPH scavenging activity of different extracts from *Viburnum grandiflorum* leaves. Each experiment was carried out in triplicates and data was shown as mean \pm SD.

***H₂O₂* reduction activity**

Further, the antioxidant activity of the extracts was determined against hydrogen peroxide. The results showed strong and dose-dependent inhibition of H_2O_2 by these four extracts from *Viburnum grandiflorum* leaves. It was observed that ethanolic extract induced strong inhibition of H_2O_2 with respect to that of reference control ascorbic acid. The ethanolic extract exhibited 40%, 51%, 68%, 76% and 83% inhibition of H_2O_2 at different concentrations ranging from 30-480 μ g/ml, respectively (Figure 2). Therefore, it was observed that the extracts from the leaves of *Viburnum grandiflorum* possess strong ability to reduce H_2O_2 radicals.




Figure 2: The hydrogen peroxide reduction activity and relative reduction by the extracts from *Viburnum grandiflorum*. Each experiment was carried out in triplicates and data was shown as mean \pm SD.

Antibacterial activity of leaf extracts of *Viburnum grandiflorum*

Medicinal plants are highly active against disease causing pathogens like bacteria, fungi, viruses etc. [20,21]. Several medicinal plants are still being used to bacterial infections in humans in different system of medicines including Ayurveda, Unani and Traditional Chinese Medicine [22]. The antibacterial activity of the extracts from leaves of *Viburnum grandiflorum* was monitored using Agar-well diffusion method. The microbes used were *M. luteus*, *S. aureus*, and *E. coli*. Each solvent extract showed significant inhibition zone against the tested bacterial strains. The highest zone of inhibition was recorded for ethanolic extract against the *M. luteus* with 30.5 ± 0.27 mm and MIC value of 0.07 in comparison to that of reference control ampicillin 34.20 ± 0.27 and MIC value of 0.01 mg/ml. The ethanolic extract was strongly active against all the three tested bacterial strains followed by ethyl acetate and n-hexane respectively. No activity was reported for the aqueous extract. The antibacterial activity of the extracts tested against selected bacterial strains is listed in table 1.

Table 1. Antibacterial activity of *Viburnum grandiflorum* leaf extracts (MIC value expressed in mg/ml).

Extract	<i>E. coli</i>		<i>S. aureus</i>		<i>M. luteus</i>	
	Inhibition zone (mm)	MIC	Inhibition zone (mm)	MIC	Inhibition zone (mm)	MIC
Ampicillin	30.9 ± 0.16	0.04	39.43 ± 0.22	0.02	34.20 ± 0.27	0.01
Ethanollic	21.7 ± 0.37	0.1	29.5 ± 0.45	0.037	30.5 ± 0.21	0.07
Ethyl acetate	19.4 ± 0.21	0.53	27.4 ± 0.31	0.07	25.3 ± 0.11	0.13
n-hexane	11.3 ± 0.17	0.91	13.4 ± 0.20	0.25	14.7 ± 0.25	0.16
Water	NA	ND	NA	ND	NA	ND

Each value is a mean of three biological replicas ± SD, NA=No activity, ND=Not determined, Ampicillin used as positive control

Cytotoxic activity of leaf extracts of *Viburnum grandiflorum*

Anticancer agents are most wanted chemical species in the world during these test and hard times of cancer resistance to conventional chemotherapy. Medicinal plants manufacture chemical species have strong activity profiles against a variety of human disorders [23]. Natural products have a remarkable potential to be a pool of anticancer research due to structural diversity and more than enough yet to be explored [24].

The cytotoxic analysis of the extracts from *Viburnum grandiflorum* leaves was evaluated via MTT assay. Results showed that the leaf extracts possess strong cytotoxicity against the lung cancer A549 cell line in a concentration-dependent manner (Figure 4). The ethanolic extract showed higher antiproliferative effects than the ethyl acetate, aqueous and n-hexane extracts. After the treatment with different concentrations ranging from 0-480 µg/ml of the ethanolic, ethyl acetate, aqueous and n-hexane extracts, the viability significantly reduced to 5%, 15%, 17% and 33%, respectively. Therefore, it was concluded that all the leaf extracts showed promising cytotoxicity against the lung cancer A549 cell line but the leading inhibitor as ethanolic extract.




Figure 3: The MTT assay results showing the cytotoxicity of different solvent extracts of *Viburnum grandiflorum* leaves. Results showed concentration-dependent inhibition of the viability by the extracts and higher activity was reported for ethanolic extract. Each experiment was carried out in triplicates and data was shown as mean ± SD.

CONCLUSION

The current research indicated that *Viburnum grandiflorum* leaves have significant health protective effects. We found that *Viburnum grandiflorum* leaf extracts showed remarkable *in vitro* cytotoxic, antimicrobial and antioxidant activities with ethanolic extract the leading one. Therefore, further investigational researches are recommended to further characterize the medicinal activities of the plant.

ACKNOWLEDGEMENT

Authors are highly thankful to the Department of Botany and Department of Pharmacy, Madhyanchal Professional University, Ratibad, Bhopal, M.P.

AUTHOR CONTRIBUTION

All the authors contributed equally in this research.

CONFLICT OF INTEREST

Authors declare that there is no conflict of interest to mention.

REFERENCES

1. IqbalAwanZahidIqbalAwan Z (2013). Ethnobotanical importance of some highly medicinal plants of district Muzaffarabad, Pakistan with special reference to the Species of the Genus Viburnum. IOSR J Pharm BiolSci 6:53-66.
2. Konarska A (2017). Comparative micromorphology and anatomy of flowers and floral secretory structures in two Viburnum species. Protoplasma 254(1):523-537.
3. Shah Z., Ali F., Ullah H., Khan D., Khan S., Khan R., Ali I. (2014). Biological Screening and Chemical Constituents of *Viburnum grandiflorum*. J. Chem. Soc. Pak, 36, 113.
4. Kumar M., Paul Y., Anand V. (2009). An ethnobotanical study of medicinal plants used by the locals in Kishtwar, Jammu and Kashmir, India. Ethnobot. Leaflet. 10, 5, 200.
5. DAR M. (2003). Ethno botanical uses of Plants of Lawat District Muzaffarabad, Azad Jammu and Kashmir. Asian J. Plant Sci. 2 (9), 680.
6. Nautiyal S., Maikhuri R., Rao K., Saxena K. (2001). Medicinal plant resources in Nanda Devi Biosphere Reserve in the central Himalayas. J. Herbs, Spices Med. Plants 8 (4), 47.
7. Latif A., Shinwari Z., Hussain J., Murtaza S. (2006). NTFPS: An alternative to forest logging in Minadam and Sultanar Valley Swat. Lyonia 11, 15.
8. Khan Z., Khuroo A., Dar G. (2004). Ethnomedicinal survey of Uri, Kashmir Himalaya. Indian J. Tradit. Know. 3 (4), 351.
9. Uddin G., Alam M., Muhammad N., Siddiqui B.S., Sadat A. (2013). Bioassay-guided Isolation of a Antinociceptive, Anti-inflammatory and Antipyretic Benzofuran Derivative from *Viburnum grandiflorum*. Pharm. Anal. Act. Vol 11, 20-26.
10. Balangcod T.D., Balangcod A.K.D. Ethnomedicinal knowledge of plants and healthcare practices among the Kalanguya tribe in Tinoc, Ifugao, Luzon, Philippines. Indian J. Tradit. Know. 10 (2), 227, 2011.
11. Alam M., Uddin G., Sadat A., Muhammad N., Khan A.A., Siddiqui B.S. Evaluation of *Viburnum grandiflorum* for its in-vitro pharmacological screening. Afr. J. Pharm. Pharmacol. 6 (22), 1606, 2012.
12. Szczepańska N., Owczarek K., Kudlak B., Pokrywka A., Mazerska Z., Gałuszka A., Namieśnik J. Analysis and Bioanalysis: an Effective Tool for Data Collection of Environmental Conditions and Processes. Pol. J. Environ. Stud. 25 (1), 45, 2016.
13. Siegel RL, Miller KD, Jemal A. (2020). Cancer statistics, CA Cancer J Clin. 2020;70(1):7-30.
14. ACSFACCS. (2020). American Cancer Society. Cancer Facts & Figures. Atlanta: American Cancer Society; 2020.
15. Majeed U, Manochakian R, Zhao Y, Lou Y.(2021).Targeted therapy in advanced non-small cell lung cancer: current advances and future trends. Journal of Hematology & Oncology. ;14(1):1-20.
16. Khursheed, A., & Jain, V. (2021). Oxidative stress-mediated caspase-dependent intrinsic apoptosis and proautophagic therapeutic potential of *Salvia moorcroftiana* Wall. exBenth. growing in the Kashmir Valley, India. Biomedical Research and Therapy, 8(11), 4718-4727.
17. Khursheed A, Jain V. (2021). Autophagy mediated anticancer activity, antioxidant and antimicrobial potential of extracts from young twigs and leaves of *Juglans regia* growing in Kashmir Valley, India. Bulletin of Environment, Pharmacology and Life Sciences, 10(6): 258-265.
18. Nasri H, Rafeian-Kopaei M. (2014). Medicinal plants and antioxidants: why they are not always beneficial?. Iranian journal of public health. ;43(2):255.
19. Keshari AK, Verma AK, Kumar T, Srivastava R. (2015). Oxidative stress: a review. The International Journal of Science and Technology. 1;3(7):155.
20. Arora DS, Kaur GJ. (2007). Antibacterial activity of some Indian medicinal plants. Journal of natural medicines. ;61(3):313-7.
21. Reshi IA, Sarkar TK, Bhat AH, Malik HU, Muhee A, Qureshi S. (2017). Antibacterial activity of some medicinal plants of Kashmir, J&K, India. Journal of Pure and Applied Microbiology. 1;11(1):521-7.
22. Tambekar DH, Dahikar SB. (2011). Antibacterial activity of some Indian Ayurvedic preparations against enteric bacterial pathogens. Journal of advanced pharmaceutical technology & research.;2(1):24.
23. Barbieri R, Coppo E, Marchese A, Daglia M, Sobarzo-Sánchez E, Nabavi SF, Nabavi SM. (2017). Phytochemicals for human disease: An update on plant-derived compounds antibacterial activity. Microbiological research. ;196:44-68.
24. Khursheed A, Jain V. (2020). Medicinal Research Progress of Natural Coumarin and its Derivatives”, The Natural Products Journal 2020;11(5):648-662.

CITATION OF THIS ARTICLE

K Trivedi and S G Singh. The Effect of phosphate Solubilizing fungus (*Aspergillus niger*) on Mustard crop productivity.Bull. Env.Pharmacol. Life Sci., Vol 11 [8] July 2022 : 40-42